SANTA CRUZ BIOTECHNOLOGY, INC.

DNA pol ε B (B-3): sc-376703



BACKGROUND

DNA replication, recombination and repair, all of which are necessary for genome stability, require the presence of exonucleases. In DNA replication, these enzymes are involved in the processing of Okazaki fragments, whereas in DNA repair, they function to excise damaged DNA fragments and correct recombinational mismatches. Exonucleases involved in these processes include DNA polymerases, including DNA pol δ and ϵ . DNA pol δ consists of two subunits, p125 which interacts directly with the sliding DNA clamp protein PCNA, and p50. DNA pol δ can be regulated by cell cycle proteins. DNA pol ϵ is a multiple subunit enzyme, the catalytic subunit of which is encoded by the POL2 gene. The exact reactions catalyzed by DNA pol δ and ϵ on leading and lagging strands have not yet been elucidated.

REFERENCES

- 1. Lee, M.Y., et al. 1984. Further studies on calf thymus DNA polymerase δ purified to homogeneity by a new procedure. Biochemistry 23: 1906-1913.
- Hamatake, R.K., et al. 1990. Purification and characterization of DNA polymerase II from the yeast *Saccharomyces cerevisiae*. Identification of the catalytic core and a possible holoenzyme form of the enzyme. J. Biol. Chem. 265: 4072-4083.
- Morrison, A., et al. 1990. A third essential DNA polymerase in *S. cerevisiae*. Cell 62: 1143-1151.
- Goulian, M., et al. 1990. Discontinuous DNA synthesis by purified mammalian proteins. J. Biol. Chem. 265: 18461-18471.
- 5. Zeng, X.R., et al. 1994. Regulation of human DNA polymerase δ during the cell cycle. J. Biol. Chem. 269: 24027-24033.
- Johnson, R.E., et al. 1995. Requirement of the yeast RTH1 5' to 3' exonuclease for the stability of simple repetitive DNA. Science 269: 238-240.
- 7. Zhang, P., et al. 1999. Direct interaction of proliferating cell nuclear antigen with the p125 catalytic subunit of mammalian DNA polymerase δ . J. Biol. Chem. 274: 26647-26653.

CHROMOSOMAL LOCATION

Genetic locus: POLE2 (human) mapping to 14q21.3; Pole2 (mouse) mapping to 12 C2.

SOURCE

DNA pol ϵ B (B-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 501-537 near the C-terminus of DNA pol ϵ B of human origin.

PRODUCT

Each vial contains 200 μg lgG_3 kappa light chain in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-376703 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

DNA pol ε B (B-3) is recommended for detection of DNA pol ε B of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DNA pol ϵ B (B-3) is also recommended for detection of DNA pol ϵ B in additional species, including equine, canine and avian.

Suitable for use as control antibody for DNA pol ϵ B siRNA (h): sc-37781, DNA pol ϵ B siRNA (m): sc-37782, DNA pol ϵ B shRNA Plasmid (h): sc-37781-SH, DNA pol ϵ B shRNA Plasmid (m): sc-37782-SH, DNA pol ϵ B shRNA (h) Lentiviral Particles: sc-37781-V and DNA pol ϵ B shRNA (m) Lentiviral Particles: sc-37782-V.

Molecular Weight of DNA pol ϵ B: 59 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, Jurkat whole cell lysate: sc-2204 or HeLa nuclear extract: sc-2120.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



DNA pol ϵ B (B-3): sc-376703. Western blot analysis of DNA pol ϵ B expression in HeLa nuclear extract.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.