

17 β -HSD (A-5): sc-376719

BACKGROUND

17 β -hydroxysteroid dehydrogenase type 1 (17 β -HSD) catalyzes the final step in the formation of estradiol and testosterone from estrone and androstenedione, respectively. Ovarian granulosa cells and breast tissue both express 17 β -HSD. Other tissues that express 17 β -HSD include testis, placenta, uterus, prostate and adipose tissue. 17 β -HSD functions as a homodimer and prefers NADP(H) over NAD(H) for oxidation and reduction. The gene encoding human 17 β -HSD maps to chromosome 17q21.2. The importance of 17 β -HSD to estradiol production suggests the specific inhibition of 17 β -HSD may aid in breast cancer therapy. Breast cancer patients with an amplification of 17 β -HSD expression statistically have a worse outcome than those without. 17 β -HSD amplification in tamoxifen-treated patients correlates to decreased breast cancer survival.

REFERENCES

1. Winqvist, R., et al. 1990. The gene for 17 β -hydroxysteroid dehydrogenase maps to human chromosome 17, bands q12-q21, and shows an RFLP with Scal. Hum. Genet. 85: 473-476.
2. Luu-The, V., et al. 1990. Structure of two in tandem human 17 β -hydroxysteroid dehydrogenase genes. Mol. Endocrinol. 4: 268-275.

CHROMOSOMAL LOCATION

Genetic locus: HSD17B1 (human) mapping to 17q21.2.

SOURCE

17 β -HSD (A-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 265-299 near the C-terminus of 17 β -HSD of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-376719 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

17 β -HSD (A-5) is recommended for detection of 17 β -HSD of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 17 β -HSD siRNA (h): sc-41381, 17 β -HSD shRNA Plasmid (h): sc-41381-SH and 17 β -HSD shRNA (h) Lentiviral Particles: sc-41381-V.


Molecular Weight of 17 β -HSD: 35 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, human placenta extract: sc-363772 or BT-20 cell lysate: sc-2223.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



17 β -HSD (A-5): sc-376719. Western blot analysis of 17 β -HSD expression in BT-20 whole cell lysate (A) and human placenta tissue extract (B).

SELECT PRODUCT CITATIONS

1. Oktem, O., et al. 2017. FSH stimulation promotes progesterone synthesis and output from human granulosa cells without luteinization. Hum. Reprod. 32: 643-652.
2. Abdel-Maksoud, F.M., et al. 2019. Prenatal exposures to bisphenol A and di (2-ethylhexyl) phthalate disrupted seminiferous tubular development in growing male rats. Reprod. Toxicol. 88: 85-90.
3. Bildik, G., et al. 2020. hCG improves luteal function and promotes progesterone output through the activation of JNK pathway in the luteal granulosa cells of the stimulated IVF cycles. Biol. Reprod. 102: 1270-1280.
4. Bildik, G., et al. 2020. Terminal differentiation of human granulosa cells as luteinization is reversed by activin-A through silencing of Jnk pathway. Cell Death Discov. 6: 93.
5. Jeminiwa, B.O., et al. 2021. Gonadal sex steroid hormone secretion after exposure of male rats to estrogenic chemicals and their combinations. Mol. Cell. Endocrinol. 533: 111332.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.