

# AMPK $\beta$ 2 (F-8): sc-376897

## BACKGROUND

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic  $\alpha$  subunit and regulatory  $\beta$  and  $\gamma$  subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate *in vivo* hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK $\alpha$ 1 and AMPK $\alpha$ 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK $\beta$ 1 encodes a 271 amino acid protein and human AMPK $\beta$ 2 encodes a 272 amino acid protein. The human AMPK $\gamma$ 1 gene encodes a 331 amino acid protein. Human AMPK $\gamma$ 2 and AMPK $\gamma$ 3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

## REFERENCES

1. Stapleton, D., et al. 1996. Mammalian AMP-activated protein kinase subfamily. *J. Biol. Chem.* 271: 611-614.
2. Stapleton, D., et al. 1997. AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409: 452-456.
3. Hardie, D.G., et al. 1997. The AMP-activated protein kinase—fuel gauge of the mammalian cell? *Eur. J. Biochem.* 246: 259-273.
4. Thornton, C., et al. 1998. Identification of a novel AMP-activated protein kinase  $\beta$  subunit isoform that is highly expressed in skeletal muscle. *J. Biol. Chem.* 273: 12443-12450.
5. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Cheung, P.C., et al. 2000. Characterization of AMP-activated protein kinase  $\gamma$  subunit isoforms and their role in AMP binding. *Biochem. J.* 346: 659-669.

## CHROMOSOMAL LOCATION

Genetic locus: PRKAB2 (human) mapping to 1q21.1; Prkab2 (mouse) mapping to 3 F2.2.

## SOURCE

AMPK $\beta$ 2 (F-8) is a mouse monoclonal antibody raised against amino acids 1-75 mapping at the N-terminus of AMPK $\beta$ 2 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgA kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

AMPK $\beta$ 2 (F-8) is recommended for detection of AMPK $\beta$ 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for AMPK $\beta$ 2 siRNA (h): sc-38927, AMPK $\beta$ 2 siRNA (m): sc-38928, AMPK $\beta$ 2 shRNA Plasmid (h): sc-38927-SH, AMPK $\beta$ 2 shRNA Plasmid (m): sc-38928-SH, AMPK $\beta$ 2 shRNA (h) Lentiviral Particles: sc-38927-V and AMPK $\beta$ 2 shRNA (m) Lentiviral Particles: sc-38928-V.


Molecular Weight of AMPK $\beta$ 2: 30-34 kDa.

Positive Controls: RT-4 whole cell lysate: sc-364257, human skeletal muscle extract: sc-363776 or A-375 cell lysate: sc-3811.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



AMPK $\beta$ 2 (F-8): sc-376897. Western blot analysis of AMPK $\beta$ 2 expression in human skeletal muscle tissue extract (A) and A-375 (B) and RT-4 (C) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.