

# NFRκB (H-5): sc-376945

## BACKGROUND

NFκB (nuclear factor κ-B) is a ubiquitously expressed transcriptional regulator that, when stimulated, can activate transcription of several genes encoding proteins involved in cell cycle control, cell adhesion and programmed cell death. NFRκB (nuclear factor related to κ-B-binding protein), also known as DNA-binding protein R κ-B, is a nuclear protein that binds to the DNA consensus sequence 5'-GGGGAATCTCC-3' of NFκB. Binding of NFRκB is thought to regulate IL-2Rα (interleukin-2 receptor α-chain) gene expression, a critical step in T-cell activation. NFRκB exists as three isoforms due to alternative splicing and is expressed primarily in the brain, liver, spleen, testis and thymus. NFRκB gene expression is amplified in acute myeloid leukemia, suggesting a possible role in carcinogenesis.

## REFERENCES

- Adams, B.S., et al. 1992. Localization of the gene encoding R κ B (NFRKB), a tissue-specific DNA binding protein, to chromosome 11q24-q25. *Genomics* 14: 270-274.
- Adams, B.S., et al. 1992. Cloning of R κ B, a novel DNA-binding protein that recognizes the interleukin-2 receptor α chain κ B site. *New Biol.* 3: 1063-1073.
- Crossen, P.E., et al. 1999. Identification of amplified genes in a patient with acute myeloid leukemia and double minute chromosomes. *Cancer Genet. Cytogenet.* 113: 126-133.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 164013. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Tyybäkinoja, A., et al. 2006. Amplified, lost, and fused genes in 11q23-25 amplicon in acute myeloid leukemia, an array-CGH study. *Genes Chromosomes Cancer* 45: 257-264.
- Natarajan, M., et al. 2006. Nuclear translocation and DNA-binding activity of NFKB (NFκB) after exposure of human monocytes to pulsed ultra-wide-band electromagnetic fields (1 kV/cm) fails to transactivate κB-dependent gene expression. *Radiat. Res.* 165: 645-654.

## CHROMOSOMAL LOCATION

Genetic locus: NFRKB (human) mapping to 11q24.3.

## SOURCE

NFRκB (H-5) is a mouse monoclonal antibody raised against amino acids 388-657 mapping within an internal region of NFRκB of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-376945 X, 200 µg/0.1 ml.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

NFRκB (H-5) is recommended for detection of NFRκB of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NFRκB (H-5) is also recommended for detection of NFRκB in additional species, including equine and canine.

Suitable for use as control antibody for NFRκB siRNA (h): sc-96360, NFRκB shRNA Plasmid (h): sc-96360-SH and NFRκB shRNA (h) Lentiviral Particles: sc-96360-V.

NFRκB (H-5) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of NFRκB: 139 kDa.

Positive Controls: IMR-32 nuclear extract: sc-2148.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



NFRκB (H-5): sc-376945. Western blot analysis of NFRκB expression in IMR-32 nuclear extract.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.