

Cdc27 (B-4): sc-376964

BACKGROUND

Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by the proteolysis of cyclins. The cell division cycle (Cdc) genes are required at various points in the cell cycle. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. Cdc6 is the human homolog of *Saccharomyces cerevisiae* Cdc6, which is involved in the initiation of DNA replication. Cdc37 appears to facilitate Cdk4/cyclin D1 complex formation and has been shown to form a stable complex with Hsp90. Cdc34, Cdc27 and Cdc16 function as ubiquitin-conjugating enzymes. Cdc34 is thought to be the structural and functional homolog of *Saccharomyces cerevisiae* Cdc34, which is essential for the G₁ to S phase transition. Cdc16 and Cdc27 are components of the APC (anaphase-promoting complex) which ubiquitinates cyclin B, resulting in cyclin B/Cdk complex degradation.

REFERENCES

- Palmer, R.E., et al. 1990. Mitotic transmission of artificial chromosomes in cdc mutants of the yeast, *Saccharomyces cerevisiae*. *Genetics* 125: 763-774.
- Gautier, J., et al. 1991. Cdc25 is a specific tyrosine phosphatase that directly activates p34^{cdc2}. *Cell* 67: 197-211.
- Plon, S.E., et al. 1993. Cloning of the human homolog of the Cdc34 cell cycle gene by complementation in yeast. *Proc. Natl. Acad. Sci. USA* 90: 10484-10488.
- King, R.W., et al. 1995. A 20S complex containing Cdc27 and Cdc16 catalyzes the mitosis-specific conjugation of ubiquitin to cyclin B. *Cell* 81: 279-288.
- Barinaga, M. 1995. A new twist to the cell cycle. *Science* 269: 631-632.

CHROMOSOMAL LOCATION

Genetic locus: CDC27 (human) mapping to 17q21.32; Cdc27 (mouse) mapping to 11 E1.

SOURCE

Cdc27 (B-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-29 at the N-terminus of Cdc27 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-376964 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Cdc27 (B-4) is recommended for detection of Cdc27 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Cdc27 (B-4) is also recommended for detection of Cdc27 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Cdc27 siRNA (h): sc-77362, Cdc27 siRNA (m): sc-35041, Cdc27 shRNA Plasmid (h): sc-77362-SH, Cdc27 shRNA Plasmid (m): sc-35041-SH, Cdc27 shRNA (h) Lentiviral Particles: sc-77362-V and Cdc27 shRNA (m) Lentiviral Particles: sc-35041-V.

Molecular Weight of Cdc27: 97 kDa.


Positive Controls: A-431 nuclear extract: sc-2122, RAW 264.7 whole cell lysate: sc-2211 or HeLa nuclear extract: sc-2120.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Cdc27 (B-4): sc-376964. Western blot analysis of Cdc27 expression in K-562 (A), MOLT-4 (B), RAW 264.7 (C) and A-10 (D) whole cell lysates.



Cdc27 (B-4): sc-376964. Western blot analysis of Cdc27 expression in A-431 (A) and HeLa (B) nuclear extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **Cdc27 (AF3.1): sc-9972** for Cdc27 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.