

# HAUSP (G-12): sc-377147

## BACKGROUND

HAUSP (herpesvirus-associated ubiquitin-specific protease, USP7) is a ubiquitin-specific protease. HAUSP localizes predominantly to the nucleus, in a TD-dependent manner, where it associates with ND10. ND10 are small nuclear structures implicated in a variety of cellular processes including response to stress and interferons, oncogenesis, and viral infection. HAUSP binds strongly to Vmw110, a herpesvirus regulatory protein which has the ability to disrupt ND10. HAUSP, a novel p53-interacting protein, functions to deubiquitinate p53 in an important pathway for p53 stabilization. HAUSP strongly stabilizes p53 even in the presence of excess Mdm2, and also induces p53-dependent cell growth repression and apoptosis. The HAUSP protein is distributed in the nucleus in a micropunctate pattern with a limited number of larger discrete foci, some of which co-localize with PML in ND10. The gene encoding HAUSP maps to human chromosome band 16p13.2.

## CHROMOSOMAL LOCATION

Genetic locus: USP7 (human) mapping to 16p13.2; Usp7 (mouse) mapping to 16 A1.

## SOURCE

HAUSP (G-12) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 17-53 near the N-terminus of HAUSP of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>3</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-377147 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

HAUSP (G-12) is recommended for detection of HAUSP of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HAUSP siRNA (h): sc-41521, HAUSP siRNA (m): sc-77373, HAUSP shRNA Plasmid (h): sc-41521-SH, HAUSP shRNA Plasmid (m): sc-77373-SH, HAUSP shRNA (h) Lentiviral Particles: sc-41521-V and HAUSP shRNA (m) Lentiviral Particles: sc-77373-V.


Molecular Weight of HAUSP: 135 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or HAUSP (m): 293T Lysate: sc-178723.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



HAUSP (G-12): sc-377147. Western blot analysis of HAUSP expression in non-transfected 293T: sc-117752 (A), mouse HAUSP transfected 293T: sc-178723 (B) and HeLa (C) whole cell lysates.



HAUSP (G-12): sc-377147. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear and cytoplasmic staining of cells in seminiferous ducts and Leydig cells.

## SELECT PRODUCT CITATIONS

- Das, S., et al. 2020. HAUSP stabilizes Cdc25A and protects cervical cancer cells from DNA damage response. *Biochim. Biophys. Acta Mol. Cell Res.* 1867: 118835.
- Gao, L., et al. 2021. Proteome analysis of USP7 substrates revealed its role in melanoma through PI3K/Akt/FOXO and AMPK pathways. *Front. Oncol.* 11: 650165.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.