

PARP-14 (C-1): sc-377150

BACKGROUND

Poly(ADP-ribosylation) is a method of DNA damage-dependent posttranslational modification that helps to rescue injured proliferating cells from cell death. The PARP (poly(ADP-ribose) polymerase) proteins comprise a superfamily of enzymes that functionally modify histones and other nuclear proteins, thereby preventing cell death. PARPs use NAD⁺ as a substrate to catalytically transfer ADP-ribose residues onto protein acceptors; a process that, when repeated multiple times, leads to the formation of poly(ADP-ribose) chains on the protein. The presence of these chains alters the function of the target protein and promotes cell survival. PARP proteins are implicated in a variety of diseases, including cancer, neurodegenerative and inflammatory disorders.

CHROMOSOMAL LOCATION

Genetic locus: PARP14 (human) mapping to 3q21.1; Parp14 (mouse) mapping to 16 B3.

SOURCE

PARP-14 (C-1) is a mouse monoclonal antibody raised against amino acids 291-358 mapping within an internal region of PARP-14 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-377150 X, 200 µg/0.1 ml.

PARP-14 (C-1) is available conjugated to agarose (sc-377150 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-377150 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-377150 PE), fluorescein (sc-377150 FITC), Alexa Fluor[®] 488 (sc-377150 AF488), Alexa Fluor[®] 546 (sc-377150 AF546), Alexa Fluor[®] 594 (sc-377150 AF594) or Alexa Fluor[®] 647 (sc-377150 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-377150 AF680) or Alexa Fluor[®] 790 (sc-377150 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

PARP-14 (C-1) is recommended for detection of PARP-14 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PARP-14 siRNA (h): sc-76056, PARP-14 siRNA (m): sc-76057, PARP-14 shRNA Plasmid (h): sc-76056-SH, PARP-14 shRNA Plasmid (m): sc-76057-SH, PARP-14 shRNA (h) Lentiviral Particles: sc-76056-V and PARP-14 shRNA (m) Lentiviral Particles: sc-76057-V.


PARP-14 (C-1) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Positive Controls: U-251-MG whole cell lysate: sc-364176, RAW 264.7 whole cell lysate: sc-2211 or MOLT-4 cell lysate: sc-2233.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



PARP-14 (C-1): sc-377150. Western blot analysis of PARP-14 expression in U-251-MG (A), MOLT-4 (B) and RAW 264.7 (C) whole cell lysates.



PARP-14 (C-1): sc-377150. Immunofluorescence staining of formalin-fixed A-431 cells showing cytoplasmic and nuclear localization (A). PARP-14 (C-1) Alexa Fluor[®] 488: sc-377150 AF488. Direct immunofluorescence staining of formalin-fixed SW480 cells showing cytoplasmic and nuclear localization. Blocked with UltraCruz[®] Blocking Reagent: sc-516214 (B).

SELECT PRODUCT CITATIONS

- Nicolae, C.M., et al. 2015. A novel role for the mono-ADP-ribosyltransferase PARP-14/ARTD8 in promoting homologous recombination and protecting against replication stress. *Nucleic Acids Res.* 43: 3143-3153.
- Becker, A.C., et al. 2018. Influenza A virus induces autophagosomal targeting of ribosomal proteins. *Mol. Cell. Proteomics* 17: 1909-1921.
- Higashi, H., et al. 2019. A study into the ADP-ribosylome of IFN-γ-stimulated THP-1 human macrophage-like cells identifies ARTD8/PARP-14 and ARTD9/PARP-9 ADP-ribosylation. *J. Proteome Res.* 18: 1607-1622.
- Dhoonmoon, A., et al. 2020. Genome-wide CRISPR synthetic lethality screen identifies a role for the ADP-ribosyltransferase PARP-14 in DNA replication dynamics controlled by ATR. *Nucleic Acids Res.* 48: 7252-7264.
- Kamata, T., et al. 2021. Post-transcriptional regulation of PARP-7 protein stability is controlled by androgen signaling. *Cells* 10: 363.
- Kuraoka, S., et al. 2022. A novel spectral annotation strategy streamlines reporting of mono-ADP-ribosylated peptides derived from mouse liver and spleen in response to IFN-γ. *Mol. Cell. Proteomics* 21: 100153.
- Chen, Q., et al. 2022. Truncated PARP-1 mediates ADP-ribosylation of RNA polymerase III for apoptosis. *Cell Discov.* 8: 3.
- Dhoonmoon, A., et al. 2022. The KU-PARP14 axis differentially regulates DNA resection at stalled replication forks by MRE11 and EXO1. *Nat. Commun.* 13: 5063.
- Xu, A.H., et al. 2023. Poly(ADP-ribose) polymerase family member 14 promotes functional recovery after spinal cord injury through regulating microglia M1/M2 polarization via STAT1/6 pathway. *Neural Regen. Res.* 18: 1809-1817.

RESEARCH USE

For research use only, not for use in diagnostic procedures.