CAPS-1 (G-12): sc-377279



The Power to Question

BACKGROUND

Calcium-dependent secretion activators (CAPS-1 and CAPS-2) are calcium-binding proteins that direct neurotransmitter and neuropeptide-filled vesicles to the cell membrane for secretory granule exocytosis. Both CAPS-1 and CAPS-2 are expressed primarily in the brain where they regulate the secretion of various substances. The CAPS proteins contain a PH domain that is essential for regulation of exocytosis, as well as regulation of phospholipid binding. Through their regulation of neurotrophin release from granule cells, CAPS proteins help to regulate cell fate during neuronal development. CAPS-1 is thought to regulate catecholamine release from neuronal cells, while CAPS-2 is thought to regulate release of both brain-derived neurotrophic factor (BDNF) and neurotrophin-3 (NT-3) from granule cells. Defects in the genes encoding CAPS-1 and CAPS-2 are implicated in impaired cerebral development and autism.

REFERENCES

- Cisternas, F.A., et al. 2003. Cloning and characterization of human CADPS and CADPS2, new members of the Ca²⁺-dependent activator for secretion protein family. Genomics 81: 279-291.
- Speidel, D., et al. 2003. A family of Ca²⁺-dependent activator proteins for secretion: comparative analysis of structure, expression, localization, and function. J. Biol. Chem. 278: 52802-52809.
- 3. Grishanin, R.N., et al. 2004. CAPS acts at a prefusion step in dense-core vesicle exocytosis as a PIP2 binding protein. Neuron 43: 551-562.
- Speidel, D., et al. 2005. CAPS-1 regulates catecholamine loading of large dense-core vesicles. Neuron 46: 75-88.

CHROMOSOMAL LOCATION

Genetic locus: CADPS (human) mapping to 3p14.2; Cadps (mouse) mapping to 14 A1.

SOURCE

CAPS-1 (G-12) is a mouse monoclonal antibody raised against amino acids 1001-1060 mapping within an internal region of CAPS-1 of human origin.

PRODUCT

Each vial contains 200 μg lgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CAPS-1 (G-12) is available conjugated to agarose (sc-377279 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-377279 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-377279 PE), fluorescein (sc-377279 FITC), Alexa Fluor® 488 (sc-377279 AF488), Alexa Fluor® 546 (sc-377279 AF546), Alexa Fluor® 594 (sc-377279 AF594) or Alexa Fluor® 647 (sc-377279 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-377279 AF680) or Alexa Fluor® 790 (sc-377279 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CAPS-1 (G-12) is recommended for detection of CAPS-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CAPS-1 siRNA (h): sc-62076, CAPS-1 siRNA (m): sc-62077, CAPS-1 shRNA Plasmid (h): sc-62076-SH, CAPS-1 shRNA Plasmid (m): sc-62077-SH, CAPS-1 shRNA (h) Lentiviral Particles: sc-62076-V and CAPS-1 shRNA (m) Lentiviral Particles: sc-62077-V.


Molecular Weight of CAPS-1: 153 kDa.


Positive Controls: rat brain extract: sc-2392 or mouse brain extract: sc-2253.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz* Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-lgG κ BP-FITC: sc-516140 or m-lgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz* Mounting Medium: sc-24941 or UltraCruz* Hard-set Mounting Medium: sc-359850.

DATA







CAPS-1 (G-12): sc-377279. Western blot analysis of CAPS-1 expression in rat brain tissue extract.

SELECT PRODUCT CITATIONS

- Zhou, H., et al. 2019. Structural and functional analysis of the CAPS SNARE-binding domain required for SNARE complex formation and exocytosis. Cell Rep. 26: 3347-3359.e6.
- 2. Weng, S., et al. 2020. CAPS1 suppresses tumorigenesis in cholangiocarcinoma. Dig. Dis. Sci. 65: 1053-1063.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA