

GODZ (A-10): sc-377378



The Power to Question

BACKGROUND

Golgi-specific DHHC (Asp-His-His-Cys) zinc finger protein (GODZ), also known as palmitoyltransferase ZDHHC3 or zinc finger protein 373, is a 327 amino acid protein member of the DHHC palmitoyltransferase family. Localized to the Golgi apparatus membrane, GODZ contains one DHHC-type zinc finger, which is necessary for its palmitoyltransferase activity. GODZ has been implicated in the palmitoylation and regulated trafficking of diverse substrates that function various inhibitory and excitatory synapses. Specifically, it palmitoylates the γ subunit 2 of GABA_A receptors, which leads to normal synaptic GABAergic inhibitory function. GODZ also palmitoylates glutamate receptors GRIA1 and GRIA2, which leads to their retention in Golgi. Two isoforms of GODZ exist as a result of alternative splicing events.

CHROMOSOMAL LOCATION

Genetic locus: ZDHHC3 (human) mapping to 3p21.31; Zdhc3 (mouse) mapping to 9 F4.

SOURCE

GODZ (A-10) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 297-327 within a C-terminal cytoplasmic domain of GODZ of human origin.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-377378 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

GODZ (A-10) is recommended for detection of GODZ of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GODZ (A-10) is also recommended for detection of GODZ in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GODZ siRNA (h): sc-75158, GODZ siRNA (m): sc-75159, GODZ shRNA Plasmid (h): sc-75158-SH, GODZ shRNA Plasmid (m): sc-75159-SH, GODZ shRNA (h) Lentiviral Particles: sc-75158-V and GODZ shRNA (m) Lentiviral Particles: sc-75159-V.

Molecular Weight (predicted) of GODZ: 37 kDa.


Molecular Weight (observed) of GODZ: 49 kDa.

Positive Controls: WEHI-231 whole cell lysate: sc-2213, EOC 20 whole cell lysate: sc-364187 or M1 whole cell lysate: sc-364782.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohisto-mount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



GODZ (A-10): sc-377378. Western blot analysis of GODZ expression in WEHI-231 (A), M1 (B), EOC 20 (C), H4 (D), RAW 264.7 (E) and PC-12 (F) whole cell lysates.



GODZ (A-10): sc-377378. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lower stomach tissue showing cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

1. Kilpatrick, C.L., et al. 2016. Dissociation of Golgi-associated DHHC-type zinc finger protein (GODZ)- and sertoli cell gene with a zinc finger domain- β (SERZ- β)-mediated palmitoylation by loss of function analyses in knock-out mice. *J. Biol. Chem.* 291: 27371-27386.
2. Du, K., et al. 2017. DHHC7 palmitoylates glucose transporter 4 (Glut4) and regulates Glut4 membrane translocation. *J. Biol. Chem.* 292: 2979-2991.
3. Yao, H., et al. 2019. Inhibiting PD-L1 palmitoylation enhances T-cell immune responses against tumours. *Nat. Biomed. Eng.* 3: 306-317.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.