

# p-p53 (B-3): sc-377553

## BACKGROUND

p53 is a DNA-binding, oligomerization domain- and transcription activation domain-containing tumor suppressor that upregulates growth arrest and apoptosis-related genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 polyubiquitinates p53 for proteasome targeting. p53 can assemble into tetramers in the absence of DNA, fluctuates between latent and active (DNA-binding) conformations, and is differentially activated through posttranslational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD) (amino acids 110-286) of p53 can compromise energetically favorable association with *cis* elements and are implicated in several human cancers. Phosphorylation of p53 at residue Thr 155 is mediated by the COP9 signalosome (CSN) and targets p53 to ubiquitin-26S Proteasome-dependent degradation.

## REFERENCES

- Hupp, T.R., et al. 1992. Regulation of the specific DNA binding function of p53. *Cell* 71: 875-876.
- Levine, A.J. 1997. p53, the cellular gatekeeper for growth and division. *Cell* 88: 323-331.

## CHROMOSOMAL LOCATION

Genetic locus: TP53 (human) mapping to 17p13.1.

## SOURCE

p-p53 (B-3) is a mouse monoclonal antibody raised against a short amino acid sequence containing Thr 55 phosphorylated p53 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

p-p53 (B-3) is available conjugated to agarose (sc-377553 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-377553 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-377553 PE), fluorescein (sc-377553 FITC), Alexa Fluor® 488 (sc-377553 AF488), Alexa Fluor® 546 (sc-377553 AF546), Alexa Fluor® 594 (sc-377553 AF594) or Alexa Fluor® 647 (sc-377553 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-377553 AF680) or Alexa Fluor® 790 (sc-377553 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-377553 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

p-p53 (B-3) is recommended for detection of Thr 55 phosphorylated p53 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for p53 siRNA (h): sc-29435, p53 shRNA Plasmid (h): sc-29435-SH and p53 shRNA (h) Lentiviral Particles: sc-29435-V.

Molecular Weight of p-p53: 53 kDa.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Lambda Phosphatase: sc-200312A and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Western blot analysis of p53 phosphorylation in untreated (A,D), ERK2 treated (B,E) and ERK2 and lambda protein phosphatase (sc-200312A) treated (C,F) human recombinant p53. Antibodies tested include p-p53 (B-3): sc-377553. (A,B,C) and p53 (Pab 240): sc-99 (D,E,F).



p-p53 (B-3): sc-377553. Immunoperoxidase staining of formalin fixed, paraffin-embedded human oral mucosa tissue showing nuclear and cytoplasmic staining of squamous epithelial cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing nuclear and cytoplasmic staining of epidermal cells (B).

## SELECT PRODUCT CITATIONS

- Tung, M.C., et al. 2013. Knockdown of lipocalin-2 suppresses the growth and invasion of prostate cancer cells. *Prostate* 73: 1281-1290.
- Marampon, F., et al. 2016. Vitamin D protects endothelial cells from irradiation-induced senescence and apoptosis by modulating MAPK/SirT1 axis. *J. Endocrinol. Invest.* 39: 411-422.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.