

p-gp130 (D-3): sc-377570

BACKGROUND

IL-6 activates intracellular signaling by binding to IL-6R (the IL-6 receptor), which subsequently associates with a second protein, known as gp130. The active signaling complex consists of at minimum IL-6, IL-6R and a dimer of two gp130 proteins that are linked by a disulfide bond. The second subunit of the IL-6 complex, gp130, also functions as a component of several additional receptor complexes, including leukemia inhibitory factor (LIF), oncostatin M (OSM), ciliary neurotrophic factor (CNTF) and IL-11. The major phosphorylation site of human gp130 is located immediately N-terminal to the di-leucine motif of gp130, which regulates the internalization of the receptor. Phosphorylation of this site, Ser 782, regulates cell surface expression of the receptor polypeptide.

REFERENCES

1. Yamasaki, K., et al. 1988. Cloning and expression of the human interleukin-6 (BSF-2/IFN β 2) receptor. *Science* 241: 825-828.
2. Taga, T., et al. 1989. Interleukin-6 triggers the association of its receptor with a possible signal transducer, gp130. *Cell* 58: 573-581.
3. Hibi, M., et al. 1990. Molecular cloning and expression of an IL-6 signal transducer, gp130. *Cell* 63: 1149-1157.
4. Murakami, M., et al. 1991. Critical cytoplasmic region of the interleukin-6 signal transducer gp130 is conserved in the cytokine receptor family. *Proc. Natl. Acad. Sci. USA* 88: 11349-11353.
5. Davis, S., et al. 1993. LIFR and gp130 as heterodimerizing signal transducers of the tripartite CNTF receptor. *Science* 260: 1805-1808.
6. Gibson, R.M., et al. 2000. Phosphorylation of human gp130 at Ser-782 adjacent to the Di-leucine internalization motif. Effects on expression and signaling. *J. Biol. Chem.* 275: 22574-22582.

CHROMOSOMAL LOCATION

Genetic locus: IL6ST (human) mapping to 5q11.2; Il6st (mouse) mapping to 13 D2.2.

SOURCE

p-gp130 (D-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 768-787 phosphorylated Ser 782 of gp130 of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_{2b}$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-377570 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

p-gp130 (D-3) is recommended for detection of Ser 782 phosphorylated gp130 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for gp130 siRNA (h): sc-29333, gp130 siRNA (m): sc-35502, gp130 shRNA Plasmid (h): sc-29333-SH, gp130 shRNA Plasmid (m): sc-35502-SH, gp130 shRNA (h) Lentiviral Particles: sc-29333-V and gp130 shRNA (m) Lentiviral Particles: sc-35502-V.


Molecular Weight of p-gp130: 130 kDa.

Positive Controls: c4 whole cell lysate: sc-364186 or 3T3-L1 cell lysate: sc-2243.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Lambda Phosphatase: sc-200312A and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Western blot analysis of gp130 phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) c4 whole cell lysates. Antibodies tested include p-gp130 (D-3): sc-377570 (A, B) and gp130 (M-20): sc-656 (C, D).

Western blot analysis of gp130 phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) 3T3-L1 whole cell lysates. Antibodies tested include p-gp130 (D-3): sc-377570 (A, B) and gp130 (M-20): sc-656 (C, D).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.