

goat anti-mouse IgG, F(ab')₂-PE-Cy5: sc-3799

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments such as the Fc or F(ab')₂ regions. Santa Cruz Biotechnology offers an extensive selection of F(ab')₂ specific secondary antibodies for immunohistochemistry and flow cytometry that are non-conjugated or labeled with either AP (alkaline phosphatase), fluorescein, biotin, FITC (fluorescein isothiocyanate), Texas Red®, TRITC (tetra-methyl rhodamine isothiocyanate), PE (phycoerythrin), PE-Cy5 (phycoerythrin with cyanin-5), PE-Cy7 (phycoerythrin with cyanin-7), APC (allophycocyanin), APC-Cy7 and (allophycocyanin with cyanin-7). F(ab')₂ secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat, and are recommended for reducing non-specific secondary antibody binding to Fc receptors on the cell surface.

SOURCE

goat anti-mouse IgG, F(ab')₂-PE-Cy5 is an affinity purified pre-adsorbed, F(ab')₂ fragment secondary antibody raised in goat against mouse IgG and conjugated to PE-Cy5 (phycoerythrin with cyanin-5).

PRODUCT

Each vial contains 200 µg goat IgG (pre-adsorbed with human IgG) in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

APPLICATIONS

goat anti-mouse IgG, F(ab')₂-PE-Cy5 is recommended for detection of mouse IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400), immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400) and flow cytometry (0.5-1 µg per 1 x 10⁶ cells). Recommended for use when trying to avoid non-specific secondary antibody binding to Fc receptors on cell surfaces.

RECOMMENDED SUPPORT PRODUCTS

- CrystalCruz™ Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- CrystalCruz™ Micro Slides 75 x 25 mm; 72 frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- Organo/Limonene Mount, non-toxic alternative to PermOUNT, 100 ml: sc-45087
- UltraCruz™ Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with *in situ* hybridization, 30 ml: sc-45088
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973
- EDTA, Disodium Salt, Dihydrate, chelating agent, 500 g: sc-29092


RESEARCH USE

For research use only, not for use in diagnostic procedures.


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



goat anti-mouse IgG F(ab')₂-PE-Cy5: sc-3799. Indirect FCM analysis of HuT 78 cells stained with CD45 (35-26), followed by PE-Cy5-conjugated goat anti-mouse IgG F(ab')₂: sc-3799. Black line histogram represents the isotype control, normal mouse IgG; 3877. Antibody tested: CD45 (35-26): sc-1178.



goat anti-mouse IgG F(ab')₂-PE-Cy5: sc-3799. Indirect FCM analysis of HuT 78 cells stained with Integrin alpha3 (VM-2), followed by PE-Cy5-conjugated goat anti-mouse IgG F(ab')₂: sc-3799. Black line histogram represents the isotype control, normal mouse IgG; 3877. Antibody tested: Integrin alpha3 (VM-2): sc-32237.

SELECT PRODUCT CITATIONS

- Rosner, C., et al. 2010. Erratum to: rhesus macaque MHC class I molecules show differential subcellular localizations. *Immunogenetics* 62: 409-418.
- Rosner, C., et al. 2011. Rhesus macaque inhibitory and activating KIR3D interact with Mamu-A-encoded ligands. *J. Immunol.* 186: 2156-2163.
- Buroker, N.E., et al. 2012. The adaptor-related protein complex 2, alpha 2 subunit (AP2alpha2) gene is a peroxisome proliferator-activated receptor cardiac target gene. *Protein J.* 31: 75-83.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Texas Red® is a registered trademark of Molecular Probes (6/02).