

hnRNP E1 siRNA (h): sc-38268

BACKGROUND

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription, pre-mRNA processing as well as mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP protein components are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm, such as hnRNP E1 and E2. hnRNP E1 may function in the cytoplasm as a translational regulatory protein, while hnRNP E2 stabilizes mRNA to enhance polioviral mRNA translation. hnRNP M is involved in pre-mRNA splicing and in stress-induced transient splicing arrest.

REFERENCES

1. Badolato, J., et al. 1995. Identification and characterisation of a novel human RNA-binding protein. *Gene* 166: 323-327.
2. Siomi, H., et al. 1995. A nuclear localization domain in the hnRNP A1 protein. *J. Cell Biol.* 129: 551-560.
3. Gattoni, R., et al. 1996. The human hnRNP-M proteins: structure and relation with early heat shock-induced splicing arrest and chromosome mapping. *Nucleic Acids Res.* 24: 2535-2542.
4. Ostareck, D.H., et al. 1997. mRNA silencing in erythroid differentiation: hnRNP K and hnRNP E1 regulate 15-lipoxygenase translation from the 3' end. *Cell* 89: 597-606.
5. Kim, J.H., et al. 2000. Protein-protein interaction among hnRNPs shuttling between nucleus and cytoplasm. *J. Mol. Biol.* 298: 395-405.
6. Melcak, I., et al. 2000. Nuclear pre-mRNA compartmentalization: trafficking of released transcripts to splicing factor reservoirs. *Mol. Biol. Cell* 11: 497-510.
7. Mahe, D., et al. 2000. Spatiotemporal regulation of hnRNP M and 2H9 gene expression during mouse embryonic development. *Biochim. Biophys. Acta* 1492: 414-424.

CHROMOSOMAL LOCATION

Genetic locus: PCBP1 (human) mapping to 2p13.3.

PRODUCT

hnRNP E1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see hnRNP E1 shRNA Plasmid (h): sc-38268-SH and hnRNP E1 shRNA (h) Lentiviral Particles: sc-38268-V as alternate gene silencing products.

For independent verification of hnRNP E1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-38268A, sc-38268B and sc-38268C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

hnRNP E1 siRNA (h) is recommended for the inhibition of hnRNP E1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

hnRNP E1 (E-2): sc-137249 is recommended as a control antibody for monitoring of hnRNP E1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor hnRNP E1 gene expression knockdown using RT-PCR Primer: hnRNP E1 (h)-PR: sc-38268-PR (20 μ l, 571 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Wang, H., et al. 2010. Proteomic and functional analyses reveal the potential involvement of endoplasmic reticulum stress and α -CP1 in the anticancer activities of oridonin in Hep G2 cells. *Integr. Cancer Ther.* 10: 160-170.
2. Meng, N., et al. 2019. Novel role of heterogeneous nuclear ribonucleoprotein E1 in regulation of apoptosis and autophagy by a triazole derivative in vascular endothelial cells. *Int. J. Biol. Sci.* 15: 1299-1309.

RESEARCH USE

For research use only, not for use in diagnostic procedures.