

CLEC-1 (E-1): sc-390005

BACKGROUND

The human β -glucan protein (Dectin-1) is a small, type II transmembrane receptor that enables β -glucan-dependent, nonopsonic recognition of zymosan and other yeast-derived particles by primary macrophages. Dectin-1 is the human homolog of the C-type (calcium dependent) lectin-like receptor (CLEC) family that play an important role in regulating innate immunity. CLEC-1 is a 280 amino acid single-pass type II transmembrane protein expressed in dendritic and endothelial cells. It accumulates in perinuclear compartments and requires an associated chain to reach the cell surface. CLEC-1 is involved in antigen uptake and is homologous to the natural killer (NK) cell receptors, NKG2s and CD94, that interact with major histocompatibility complex class I molecules and either inhibit or activate cytotoxicity and cytokine secretion. It has a single carbohydrate recognition domain with six conserved and two additional cysteine residues. Additionally, CLEC-1 has a cytoplasmic immunoreceptor tyrosine-based motif and many potential phosphorylation sites.

REFERENCES

- Colonna, M., Samaridis, J. and Angman, L. 2000. Molecular characterization of two novel C-type lectin-like receptors, one of which is selectively expressed in human dendritic cells. *Eur. J. Immunol.* 30: 697-704.
- Sobanov, Y., Bernreiter, A., Derdak, S., Mechtcheriakova, D., Schweighofer, B., Döchler, M., Kalthoff, F. and Hofer, E. 2001. A novel cluster of lectin-like receptor genes expressed in monocytic, dendritic and endothelial cells maps close to the NK receptor genes in the human NK gene complex. *Eur. J. Immunol.* 31: 3493-3503.
- Lu, Q., Navdaev, A., Clemetson, J.M. and Clemetson, K.J. 2005. Snake venom C-type lectins interacting with platelet receptors. Structure-function relationships and effects on haemostasis. *Toxicon* 45: 1089-1098.
- Rosen, D.B., Bettadapura, J., Alsharifi, M., Mathew, P.A., Warren, H.S. and Lanier, L.L. 2005. Cutting edge: lectin-like transcript-1 is a ligand for the inhibitory human NKR-P1A receptor. *J. Immunol.* 175: 7796-7799.
- Rupp, C., Dolznig, H., Puri, C., Sommergruber, W., Kerjaschki, D., Rettig, W.J. and Garin-Chesa, P. 2006. Mouse endosialin, a C-type lectin-like cell surface receptor: expression during embryonic development and induction in experimental cancer neoangiogenesis. *Cancer Immun.* 6: 10.
- Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 606782. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: CLEC1A (human) mapping to 12p13.2.

SOURCE

CLEC-1 (E-1) is a mouse monoclonal antibody raised against amino acids 124-224 mapping within an internal region of CLEC-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CLEC-1 (E-1) is recommended for detection of CLEC-1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CLEC-1 siRNA (h): sc-60396, CLEC-1 shRNA Plasmid (h): sc-60396-SH and CLEC-1 shRNA (h) Lentiviral Particles: sc-60396-V.


Molecular Weight of CLEC-1: 32 kDa.

Positive Controls: CLEC-1 (h): 293T Lysate: sc-114490.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CLEC-1 (E-1): sc-390005. Western blot analysis of CLEC-1 expression in non-transfected: sc-117752 (A) and human CLEC-1 transfected: sc-114490 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.