

# ADAMTS-L4 (H-9): sc-390187

## BACKGROUND

ADAMTS (a disintegrin and metalloproteinase domain with thrombospondin type 1 modules) is a family of zinc-dependent proteases that are implicated in a variety of normal and pathological conditions, including arthritis and cancer. ADAMTS protein family members contain an amino-terminal propeptide domain, a metalloproteinase domain, a disintegrin-like domain and a carboxy-terminus that contains a varying number of Thrombospondin type 1 (TSP-1) motifs. ADAMTS-L4 (ADAMTS-like protein 4), also known as TSRC1 (thrombospondin repeat-containing protein 1), is a 1,074 amino acid secreted protein. Known to interact with cathepsin B, ADAMTS-L4 functions as a positive regulator of apoptosis. Mutations in the gene that encodes ADAMTS-L4 are a cause of autosomal recessive isolated ectopia lentis (EL), a rare condition in which defective zonule formation results in partial or complete displacement of the lens from its space. ADAMTS-L4 is expressed in spleen, liver, skeletal muscle, lung, colon, testis, placenta heart and leukocyte.

## REFERENCES

- Adams, J.C. and Tucker, R.P. 2000. The thrombospondin type 1 repeat (TSR) superfamily: diverse proteins with related roles in neuronal development. *Dev. Dyn.* 218: 280-299.
- Buchner, D.A. and Meisler, M.H. 2003. TSRC1, a widely expressed gene containing seven thrombospondin type I repeats. *Gene* 307: 23-30.
- Liu, T., et al. 2005. Human plasma N-glycoproteome analysis by immunoaffinity subtraction, hydrazide chemistry, and mass spectrometry. *J. Proteome Res.* 4: 2070-2080.
- Liu, J., et al. 2006. Cathepsin B and its interacting proteins, bikunin and TSRC1, correlate with TNF-induced apoptosis of ovarian cancer cells OV-90. *FEBS Lett.* 580: 245-250.

## CHROMOSOMAL LOCATION

Genetic locus: ADAMTSL4 (human) mapping to 1q21.3; Adamtsl4 (mouse) mapping to 3 F2.1.

## SOURCE

ADAMTS-L4 (H-9) is a mouse monoclonal antibody raised against amino acids 586-835 mapping within an internal region of ADAMTS-L4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ADAMTS-L4 (H-9) is available conjugated to agarose (sc-390187 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390187 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390187 PE), fluorescein (sc-390187 FITC), Alexa Fluor® 488 (sc-390187 AF488), Alexa Fluor® 546 (sc-390187 AF546), Alexa Fluor® 594 (sc-390187 AF594) or Alexa Fluor® 647 (sc-390187 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-390187 AF680) or Alexa Fluor® 790 (sc-390187 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

ADAMTS-L4 (H-9) is recommended for detection of ADAMTS-L4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ADAMTS-L4 siRNA (h): sc-88067, ADAMTS-L4 siRNA (m): sc-140870, ADAMTS-L4 shRNA Plasmid (h): sc-88067-SH, ADAMTS-L4 shRNA Plasmid (m): sc-140870-SH, ADAMTS-L4 shRNA (h) Lentiviral Particles: sc-88067-V and ADAMTS-L4 shRNA (m) Lentiviral Particles: sc-140870-V.


Molecular Weight of ADAMTS-L4 isoforms: 116/95 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, AMJ2-C8 whole cell lysate: sc-364366 or rat liver extract: sc-2395.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



ADAMTS-L4 (H-9): sc-390187. Western blot analysis of ADAMTS-L4 expression in AMJ2-C8 (A) and HeLa (B) whole cell lysates and rat liver tissue extract (C).



ADAMTS-L4 (H-9): sc-390187. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane and cytoplasmic localization.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.