SANTA CRUZ BIOTECHNOLOGY, INC.

eIF5A (H-8): sc-390202



BACKGROUND

In mammalian cells, translation is controlled at the level of polypeptide chain initiation by eukaryotic initiation factors. The human eukaryotic translation initiation factor 5A gene, also designated eIF-4D or eIF5A1, maps to chromosome 17p131 and encodes a 154 amino acid protein that is linked to cellular polyamine homeostasis. eIF5A1 localizes to the nuclear and cytoplasmic compartments of mammalian cells where it can stimulate ribosomal peptidyltransferase and may be involved in nucleocytoplasmic mRNA transport and/or protein translation. eIF5A1 contains a unique spermidine-derived post-translational modification at Lys-50, hypusine, which is necessary for eIF5A1's biochemical activity and for cellular proliferative signaling. In addition, eIF5A1 is a cellular cofactor for the function of the Rev transactivator protein of human immunodeficiency virus type 1 (HIV-1). Inhibition of eIF5A1 interaction with Rev leads to a block of the viral replication cycle. A highly-conserved protein that is found in all living organisms, eIF5A2 (eukaryotic translation initiation factor 5A-2) is a 153 amino acid protein that has 94% sequence similarity to elF5A1 and also shares the hypusine residue. Amplification of the gene encoding eIF5A2 is observed in ovarian carcinomas and overexpression of eIF5A2 is linked to advanced stages of ovarian cancers.

SOURCE

eIF5A (H-8) is a mouse monoclonal antibody raised against amino acids 9-124 mapping near the N-terminus of eIF5A1 of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lgG_{2a}$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

eIF5A (H-8) is available conjugated to agarose (sc-390202 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390202 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390202 PE), fluorescein (sc-390202 FITC), Alexa Fluor[®] 488 (sc-390202 AF488), Alexa Fluor[®] 546 (sc-390202 AF546), Alexa Fluor[®] 594 (sc-390202 AF594) or Alexa Fluor[®] 647 (sc-390202 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-390202 AF680) or Alexa Fluor[®] 790 (sc-390202 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

elF5A (H-8) is recommended for detection of elF5A1 and elF5A2 of mouse, rat and human origin and elF5AL1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

eIF5A (H-8) is also recommended for detection of eIF5A1, eIF5A2 and eIF5AL1 in additional species, including equine, canine, bovine and porcine.

Molecular Weight of eIF5A: 17 kDa.

Positive Controls: HUV-EC-C whole cell lysate: sc-364180, Jurkat whole cell lysate: sc-2204 or SW480 cell lysate: sc-2219.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





elF5A (H-8): sc-390202. Western blot analysis of elF5A expression in A-431 (**A**), MOLT-4 (**B**), C2C12 (**C**), I-11.15 (**D**), NRK (**E**) and A-10 (**F**) whole cell lysates. eIF5A (H-8): sc-390202. Western blot analysis of eIF5A expression in HUV-EC-C (A), Jurkat (B) and SW480 (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Chen, D.J., et al. 2014. Cadmium induces cytotoxicity in human bronchial epithelial cells through upregulation of eIF5A1 and NFκB. Biochem. Biophys. Res. Commun. 445: 95-99.
- Chen, D.J., et al. 2015. Proteomic analysis of secreted proteins by human bronchial epithelial cells in response to cadmium toxicity. Proteomics 15: 3075-3086.
- Xu, Y.M., et al. 2019. Cadmium telluride quantum dot-exposed human bronchial epithelial cells: a further study of the cellular response by proteomics. Toxicol. Res. 8: 994-1001.
- Rizzotto, D., et al. 2020. Nutlin-induced apoptosis is specified by a translation program regulated by PCBP2 and DHX30. Cell Rep. 30: 4355-4369.e6.
- Balukoff, N.C., et al. 2020. A translational program that suppresses metabolism to shield the genome. Nat. Commun. 11: 5755.
- Becker, A.E., et al. 2021. eIF5A-independent role of DHPS in p21^{CIP1} and cell fate regulation. Int. J. Mol. Sci. 22: 13187.
- Ha, D.P., et al. 2022. Targeting GRP78 suppresses oncogenic KRAS protein expression and reduces viability of cancer cells bearing various KRAS mutations. Neoplasia 33: 100837.
- Park, B.S., et al. 2024. Polyamine and EIF5A hypusination downstream of c-Myc confers targeted therapy resistance in BRAF mutant melanoma. Mol. Cancer 23: 136.
- Yamamoto, V., et al. 2024. GRP78 inhibitor YUM70 upregulates 4E-BP1 and suppresses c-MYC expression and viability of oncogenic c-MYC tumors. Neoplasia 55: 101020.

RESEARCH USE

For research use only, not for use in diagnostic procedures.