

# TEX44 (H-2): sc-390593

## BACKGROUND

Chromosome 2, the second largest human chromosome, consists of 237 million bases encoding over 1,400 genes, comprising approximately 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alström syndrome is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and vestigial telomeres, which gives credence to the hypothesis that human chromosome 2 is the result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes.

## REFERENCES

1. Ijdo, J.W., et al. 1991. Origin of human chromosome 2: an ancestral telomere-telomere fusion. *Proc. Natl. Acad. Sci. USA* 88: 9051-9055.
2. Avarello, R., et al. 1992. Evidence for an ancestral alphoid domain on the long arm of human chromosome 2. *Hum. Genet.* 89: 247-249.
3. Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. *Nature* 434: 724-731.
4. Thomas, A.C., et al. 2006. ABCA12 is the major harlequin ichthyosis gene. *J. Invest. Dermatol.* 126: 2408-2413.
5. Akiyama, M., et al. 2007. Compound heterozygous ABCA12 mutations including a novel nonsense mutation underlie harlequin ichthyosis. *Dermatology* 215: 155-159.

## CHROMOSOMAL LOCATION

Genetic locus: TEX44 (human) mapping to 2q37.1; Tex44 (mouse) mapping to 1 D.

## SOURCE

TEX44 (H-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 371-397 of TEX44 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

TEX44 (H-2) is available conjugated to agarose (sc-390593 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-390593 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-390593 PE), fluorescein (sc-390593 FITC), Alexa Fluor® 488 (sc-390593 AF488), Alexa Fluor® 546 (sc-390593 AF546), Alexa Fluor® 594 (sc-390593 AF594) or Alexa Fluor® 647 (sc-390593 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-390593 AF680) or Alexa Fluor® 790 (sc-390593 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-390593 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

TEX44 (H-2) is recommended for detection of TEX44 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TEX44 siRNA (h): sc-94299, TEX44 siRNA (m): sc-108359, TEX44 shRNA Plasmid (h): sc-94299-SH, TEX44 shRNA Plasmid (m): sc-108359-SH, TEX44 shRNA (h) Lentiviral Particles: sc-94299-V and TEX44 shRNA (m) Lentiviral Particles: sc-108359-V.


Molecular Weight of TEX44: 42 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, U-251-MG whole cell lysate: sc-364176 or T98G cell lysate: sc-2294.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



TEX44 (H-2): sc-390593. Western blot analysis of TEX44 expression in U-251-MG (A), T98G (B), Neuro-2A (C) and EOC 20 (D) whole cell lysates and mouse postnatal brain (E) and rat brain (F) tissue extracts.



TEX44 (H-2): sc-390593. Western blot analysis of TEX44 expression in MCF7 (A) and U-251-MG (B) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.