# SANTA CRUZ BIOTECHNOLOGY, INC.

# donkey anti-sheep IgG-TR: sc-3913



## BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments. Santa Cruz Biotechnology offers an extensive selection of secondary antibodies optimized for immuno-histochemistry and flow cytometry, and are labeled with either biotin, FITC (fluorescein isothiocyanate), Texas Red<sup>®</sup>, TRITC (tetramethyl rhodamine iso-thiocyanate), PE (phycoerythrin), PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex with cyanin-5.5). Immuno-histochemistry and flow cytometry secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat.

## SOURCE

donkey anti-sheep IgG-TR is a pre-adsorbed, affinity purified secondary antibody raised in donkey against sheep IgG and conjugated to Texas Red<sup>®</sup>.

#### PRODUCT

Each vial contains 200  $\mu g$  donkey IgG (pre-adsorbed with mouse and human IgG) in 0.5 ml of PBS containing 0.02% sodium azide.

#### APPLICATIONS

donkey anti-sheep IgG-TR is recommended for detection of sheep IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400) and immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400).

### **RECOMMENDED SUPPORT PRODUCTS**

#### A. TISSUE CULTURE CELLS

- CrystalCruz<sup>™</sup> Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- CrystalCruz<sup>™</sup> Micro Slides 75 x 25 mm; 72 frosted sides: sc-24976
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336

#### **B. FROZEN TISSUE SECTIONS**

- Organo/Limonene Mount, non-toxic alternative to Permount, 100 ml: sc-45087
- UltraCruz™ Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with in situ hybridization, 30 ml: sc-45088

#### C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS

- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

Texas Red<sup>®</sup> is a registered trademark of Molecular Probes (6/02).

### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.