

PTEN (H-3): sc-393088

BACKGROUND

As human tumors progress to advanced stages, one genetic alteration that occurs at high frequency is a loss of heterozygosity (LOH) at chromosome 10q23.31. Mapping of homozygous deletions on this chromosome led to the isolation of the PTEN gene, also designated MMAC1 (for mutated in multiple advanced cancers) and TEP1. This candidate tumor suppressor gene exhibits a high frequency of mutations in human glioblastomas and is also mutated in other cancers, including sporadic brain, breast, kidney and prostate cancers. PTEN has been associated with Cowden disease, an autosomal dominant cancer predisposition syndrome. The PTEN gene product is a putative protein tyrosine phosphatase that is localized to the cytoplasm and shares extensive homology with the cytoskeletal proteins tensin and auxilin. Gene transfer studies have indicated that the phosphatase domain of PTEN is essential for growth suppression of glioma cells.

CHROMOSOMAL LOCATION

Genetic locus: PTEN (human) mapping to 10q23.31; Pten (mouse) mapping to 19 C1.

SOURCE

PTEN (H-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-28 at the N-terminus of PTEN of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-393088 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

PTEN (H-3) is recommended for detection of PTEN of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PTEN (H-3) is also recommended for detection of PTEN in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for PTEN siRNA (h): sc-29459, PTEN siRNA (m): sc-36326, PTEN siRNA (r): sc-61873, PTEN shRNA Plasmid (h): sc-29459-SH, PTEN shRNA Plasmid (m): sc-36326-SH, PTEN shRNA Plasmid (r): sc-61873-SH, PTEN shRNA (h) Lentiviral Particles: sc-29459-V, PTEN shRNA (m) Lentiviral Particles: sc-36326-V and PTEN shRNA (r) Lentiviral Particles: sc-61873-V.


Molecular Weight of PTEN: 55 kDa.

Positive Controls: PTEN (m): 293T Lysate: sc-122834.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



PTEN (H-3): sc-393088. Western blot analysis of PTEN expression in non-transfected: sc-117752 (A) and mouse PTEN transfected: sc-122834 (B) 293T whole cell lysates.



PTEN (H-3): sc-393088. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lower stomach tissue showing cytoplasmic staining of glandular cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing cytoplasmic and nuclear staining of keratinocytes, fibroblasts, Langerhans cells and melanocytes (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **PTEN (A2B1): sc-7974** for PTEN antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.