

FXYD3 (B-3): sc-393639



The Power to Question

BACKGROUND

The mammalian FXYD family maintains Na⁺ and K⁺ gradients between the intracellular and extracellular milieus of cells in processes such as renal Na⁺-reabsorption, muscle contraction and neuronal excitability. FXYDs are single-span membrane proteins that share a 35 amino acid signature domain, beginning with the sequence PFXVD and containing seven invariant and six conserved amino acids. Members of the FXYD family include FXYD1 (PLM, phospholemman), FXYD2 (the γ subunit of the Na/K-ATPase), FXYD3 (Mat8, mammary tumor protein), FXYD4 (CHIF) and FXYD5 (RIC). FXYD3, a 67 amino acid protein, may act as a chloride channel or as a chloride channel regulator. It is expressed in a subset of human breast tumors and shows partial homology to FXYD1. FXYD3 has a putative 20 amino acid leader sequence and a putative transmembrane domain (with two cysteine residues). It contains no consensus phosphorylation sites in the cytoplasmic domain.

REFERENCES

1. Morrison, B.W. and Leder, P. 1994. Neu and Ras initiate murine mammary tumors that share genetic markers generally absent in c-Myc and Int-2-initiated tumors. *Oncogene* 9: 3417-3426.
2. Morrison, B.W., et al. 1995. Mat8, a novel phospholemman-like protein expressed in human breast tumors, induces a chloride conductance in *Xenopus* oocytes. *J. Biol. Chem.* 270: 2176-2182.
3. Sweadner, K.J. and Rael, E. 2000. The FXYD gene family of small ion transport regulators or channels: cDNA sequence, protein signature sequence, and expression. *Genomics* 68: 41-56.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604996. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Crambert, G., et al. 2005. FXYD3 (Mat8), a new regulator of Na,K-ATPase. *Mol. Biol. Cell* 16: 2363-2371.

CHROMOSOMAL LOCATION

Genetic locus: Fxyd3 (mouse) mapping to 7 B1.

SOURCE

FXYD3 (B-3) is a mouse monoclonal antibody raised against amino acids 1-88 representing full length FXYD3 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

FXYD3 (B-3) is available conjugated to agarose (sc-393639 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393639 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393639 PE), fluorescein (sc-393639 FITC), Alexa Fluor® 488 (sc-393639 AF488), Alexa Fluor® 546 (sc-393639 AF546), Alexa Fluor® 594 (sc-393639 AF594) or Alexa Fluor® 647 (sc-393639 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-393639 AF680) or Alexa Fluor® 790 (sc-393639 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

FXYD3 (B-3) is recommended for detection of FXYD3 of mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FXYD3 siRNA (m): sc-60666, FXYD3 shRNA Plasmid (m): sc-60666-SH and FXYD3 shRNA (m) Lentiviral Particles: sc-60666-V.

Molecular Weight of FXYD3: 8 kDa.


Positive Controls: FXYD3 (m): 293T Lysate: sc-126876, rat pancreas extract: sc-364806 or mouse pancreas extract: sc-364244.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



FXYD3 (B-3): sc-393639. Western blot analysis of FXYD3 expression in non-transfected: sc-117752 (A) and mouse FXYD3 transfected: sc-126876 (B) 293T whole cell lysates and mouse pancreas tissue extract (C).



FXYD3 (B-3): sc-393639. Western blot analysis of FXYD3 expression in mouse pancreas (A) and rat pancreas (B) tissue extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA