

netrin G1 (H-4): sc-393665

BACKGROUND

Netrin G1 and netrin G2, also referred to as laminin-1 and laminin-2, are membrane bound axon guidance molecules involved in synaptic formation and maintenance. They comprise a subgroup within the UNC-6/netrin family. Both genes have been associated with schizophrenia involving single nucleotide polymorphisms. They are both expressed in the brain but G₁ is most predominantly expressed in the thalamus and G₂ is most predominantly expressed in the cortex. These two proteins differ from classical netrins by their failure to bind netrin receptors, the presence of a glycosyl phosphatidylinositol membrane anchor, and the generation of multiple isoforms. Netrin G1 has at least nine isoforms, all of which are expressed in adult brain. Isoforms G1a, c, d, and e are also expressed in fetal brain. G1c and G1d are the most highly expressed netrin G1 isoforms. Netrin G1 is involved in NMDA receptor function and may play a role in Rett syndrome (RTT), atypical autism, epilepsy and mental retardation.

REFERENCES

1. Yin, Y., et al. 2002. Laminins: laminin- and netrin-related genes expressed in distinct neuronal subsets. *Mol. Cell. Neurosci.* 19: 344-358.
2. Fukasawa, M., et al. 2004. Case-control association study of human netrin G1 gene in Japanese schizophrenia. *J. Med. Dent. Sci.* 51: 121-128.
3. Miyashita, T., et al. 2005. Strong expression of netrin G2 in the monkey claustrum. *Neuroscience* 136: 487-496.
4. Meerabux, J.M., et al. 2005. Human netrin G1 isoforms show evidence of differential expression. *Genomics* 86: 112-116.
5. Borg, I., et al. 2005. Disruption of netrin G1 by a balanced chromosome translocation in a girl with Rett syndrome. *Eur. J. Hum. Genet.* 13: 921-927.
6. Aoki-Suzuki, M., et al. 2005. A family-based association study and gene expression analyses of netrin G1 and G2 genes in schizophrenia. *Biol. Psychiatry* 57: 382-393.
7. Archer, H.L., et al. 2006. NTNG1 mutations are a rare cause of Rett syndrome. *Am. J. Med. Genet. A* 140: 691-694.
8. Bisgaard, A.M., et al. 2007. Interstitial deletion of the short arm of chromosome 1 (1p13.1-p21.1) in a girl with mental retardation, short stature and colobomata. *Clin. Dysmorphol.* 16: 109-112.
9. Eastwood, S.L. and Harrison, P.J. 2008. Decreased mRNA expression of netrin G1 and netrin G2 in the temporal lobe in schizophrenia and bipolar disorder. *Neuropsychopharmacology* 33: 933-945.

CHROMOSOMAL LOCATION

Genetic locus: NTNG1 (human) mapping to 1p13.3.

SOURCE

netrin G1 (H-4) is a mouse monoclonal antibody raised against amino acids 175-229 mapping within an internal region of netrin G1 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

netrin G1 (H-4) is recommended for detection of netrin G1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for netrin G1 siRNA (h): sc-72290, netrin G1 shRNA Plasmid (h): sc-72290-SH and netrin G1 shRNA (h) Lentiviral Particles: sc-72290-V.


Molecular Weight of netrin G1: 18 kDa.

Positive Controls: netrin G1 (h): 293T Lysate: sc-114149.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



netrin G1 (H-4): sc-393665. Western blot analysis of netrin G1 expression in non-transfected: sc-117752 (A) and human netrin G1 transfected: sc-114149 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.