

NELF-D (E-8): sc-393682

BACKGROUND

NELF-D (negative elongation factor C/D), also known as TH1, NELF-C, HSPC130 or TH1L, is a 590 amino acid protein that localizes to the nucleus and exists as a component of the multi-protein NELF complex, a structure which negatively regulates Pol II-dependent transcription elongation. Expressed in a variety of tissues, including liver, heart, kidney, lung, brain, placenta and pancreas, NELF-D is involved in controlling transcriptional pausing of Pol II and may be able to induce chromatin unfolding, possibly playing a role in tumorigenesis. NELF-D is encoded by a gene which maps to human chromosome 20. Comprising approximately 2% of the human genome, chromosome 20 contains nearly 63 million bases that encode over 600 genes, some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome. Additionally, chromosome 20 contains a region with numerous genes which are thought important for seminal production and may be potential targets for male contraception.

REFERENCES

1. Ping, Y.H. and Rana, T.M. 2001. DSIF and NELF interact with RNA polymerase II elongation complex and HIV-1 Tat stimulates P-TEFb-mediated phosphorylation of RNA polymerase II and DSIF during transcription elongation. *J. Biol. Chem.* 276: 12951-12958.
2. Narita, T., et al. 2003. Human transcription elongation factor NELF: identification of novel subunits and reconstitution of the functionally active complex. *Mol. Cell. Biol.* 23: 1863-1873.
3. Wu, C.H., et al. 2005. Molecular characterization of *Drosophila* NELF. *Nucleic Acids Res.* 33: 1269-1279.
4. Joó, J.G., et al. 2006. Trisomy 20 mosaicism and nonmosaic trisomy 20: a report of 2 cases. *J. Reprod. Med.* 51: 209-212.
5. Ville, D., et al. 2006. Early pattern of epilepsy in the ring chromosome 20 syndrome. *Epilepsia* 47: 543-549.
6. Elghezal, H., et al. 2007. Ring chromosome 20 syndrome without deletions of the subtelomeric and CHRNA4—KCNQ2 genes loci. *Eur. J. Med. Genet.* 50: 441-445.
7. Kazantsev, A.G. 2007. Cellular pathways leading to neuronal dysfunction and degeneration. *Drug News Perspect.* 20: 501-509.
8. Lundwall, A. 2007. A locus on chromosome 20 encompassing genes that are highly expressed in the epididymis. *Asian J. Androl.* 9: 540-544.
9. Zhang, Z., et al. 2007. Negative elongation factor NELF represses human immunodeficiency virus transcription by pausing the RNA polymerase II complex. *J. Biol. Chem.* 282: 16981-16988.

CHROMOSOMAL LOCATION

Genetic locus: NELFCD (human) mapping to 20q13.32; Nelfcd (mouse) mapping to 2 H4.

SOURCE

NELF-D (E-8) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of NELF-D of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NELF-D (E-8) is recommended for detection of NELF-D of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NELF-D siRNA (h): sc-75898, NELF-D siRNA (m): sc-149908, NELF-D shRNA Plasmid (h): sc-75898-SH, NELF-D shRNA Plasmid (m): sc-149908-SH, NELF-D shRNA (h) Lentiviral Particles: sc-75898-V and NELF-D shRNA (m) Lentiviral Particles: sc-149908-V.


Molecular Weight of NELF-D: 59 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or MCF7 whole cell lysate: sc-2206.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NELF-D (E-8): sc-393682. Western blot analysis of NELF-D expression in HeLa (A), MCF7 (B) and NIH/3T3 (C) whole cell lysates. Note lack of reactivity with mouse NELF-D in lane C.



NELF-D (E-8): sc-393682. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.