

8D6 (F-12): sc-393892

BACKGROUND

The 8D6 protein, also known as 8D6A, CD320 or FDC-SM-8D6, is a single pass, type I membrane protein with two low-density lipoprotein receptor ligand binding repeats (LDL-A modules). It is expressed by follicular dendritic cells in the germinal center and acts as a stimulatory signaling molecule. Follicular dendritic cells and T cells interact with germinal center B cells, providing signals for survival, proliferation and differentiation into memory B cells and plasma cells. A disruption of this interaction results in apoptosis of B cells. 8D6 is a growth factor required for plasma cell generation from germinal center B cells. Protein 8D6, together with HCAM (or CD44), plays a significant role in the proliferation of lymphoma cells of germinal center origin. 8D6 is responsible for enhancing proliferation while HCAM inhibits apoptosis.

REFERENCES

1. Shaw, M.A. 1987. Monoclonal anti-LWab and anti-D reagents recognize a number of different epitopes. Use of red cells of non-human primates. *J. Immunogenet.* 13: 377-386.
2. Itohara, S., et al. 1989. Monoclonal antibodies specific to native murine T cell receptor $\gamma\delta$: analysis of $\gamma\delta$ T cells during thymic ontogeny and in peripheral lymphoid organs. *Proc. Natl. Acad. Sci. USA* 86: 5094-5098.
3. Kirsch, A.H., et al. 1997. The pattern of expression of CD147/neurothelin during human T cell ontogeny as defined by the monoclonal antibody 8D6. *Tissue Antigens* 50: 147-152.
4. Li, L., et al. 2000. Identification of a human follicular dendritic cell molecule that stimulates germinal center B cell growth. *J. Exp. Med.* 191: 1077-1084.
5. Zhang, X., et al. 2001. The distinct roles of T cell-derived cytokines and a novel follicular dendritic cell-signaling molecule 8D6 in germinal center-B cell differentiation. *J. Immunol.* 167: 49-56.

CHROMOSOMAL LOCATION

Genetic locus: CD320 (human) mapping to 19p13.2.

SOURCE

8D6 (F-12) is a mouse monoclonal antibody raised against amino acids 152-282 mapping at the C-terminus of 8D6 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

8D6 (F-12) is available conjugated to agarose (sc-393892 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393892 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393892 PE), fluorescein (sc-393892 FITC), Alexa Fluor[®] 488 (sc-393892 AF488), Alexa Fluor[®] 546 (sc-393892 AF546), Alexa Fluor[®] 594 (sc-393892 AF594) or Alexa Fluor[®] 647 (sc-393892 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-393892 AF680) or Alexa Fluor[®] 790 (sc-393892 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

8D6 (F-12) is recommended for detection of 8D6 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 8D6 siRNA (h): sc-106884, 8D6 shRNA Plasmid (h): sc-106884-SH and 8D6 shRNA (h) Lentiviral Particles: sc-106884-V.

Molecular Weight (predicted) of 8D6: 29 kDa.


Molecular Weight (observed) of 8D6: 38 kDa.

Positive Controls: COLO 320DM cell lysate: sc-2226, SW480 cell lysate: sc-2219 or NCI-H929 whole cell lysate: sc-364786.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



8D6 (F-12): sc-393892. Western blot analysis of 8D6 expression in NCI-H929 (A), COLO 320DM (B) and U-251-MG (C) whole cell lysates.



8D6 (F-12): sc-393892. Western blot analysis of 8D6 expression in SW480 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA