

# α-lactalbumin (H-1): sc-393900

## BACKGROUND

α-lactalbumin is the B protein of lactose synthetase secreted by the mammary epithelial cells. It is a potent Ca<sup>2+</sup>-elevating and apoptosis-inducing agent with broad, yet selective, cytotoxic activity. Multimeric α-lactalbumin has been shown to kill all transformed, embryonic and lymphoid cells tested, but not mature epithelial elements. This suggests that milk contributes to mucosal immunity not only by furnishing antimicrobial molecules but also by policing the function of lymphocytes and epithelium. α-lactalbumin may be helpful in discovering the site of origin of metastatic breast tumors. Human lactalbumin contains 123 amino acid residues. Comparison of the 5' flanking sequences of the two α-lactalbumin genes with those of five casein genes reveals the presence of a highly conserved region extending from position -140 to -110 in all seven sequences examined, suggesting a possible regulatory role in the hormonal control or tissue-specific expression of milk protein genes in the mammary gland.

## REFERENCES

- Burchell, J., et al. 1985. Production and characterization of monoclonal antibodies to human casein. A monoclonal antibody that cross-reacts with casein and α-lactalbumin. *Hybridoma* 4: 341-350.
- Wang, Q., et al. 2006. Quantitative assessment of thermal denaturation of bovine α-lactalbumin via low-intensity ultrasound, HPLC, and DSC. *J. Agric. Food Chem.* 54: 6501-6506.
- Anema, S.G., et al. 2006. Effect of protein, nonprotein-soluble components, and lactose concentrations on the irreversible thermal denaturation of β-lactoglobulin and α-lactalbumin in skim milk. *J. Agric. Food Chem.* 54: 7339-7348.
- Barros, R.M., et al. 2006. Molecular characterization of peptides released from β-lactoglobulin and α-lactalbumin via cardosins A and B. *J. Dairy Sci.* 89: 483-494.

## CHROMOSOMAL LOCATION

Genetic locus: LALBA (human) mapping to 12q13.11.

## SOURCE

α-lactalbumin (H-1) is a mouse monoclonal antibody raised against amino acids 1-142 representing full length α-lactalbumin of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

α-lactalbumin (H-1) is available conjugated to agarose (sc-393900 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-393900 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-393900 PE), fluorescein (sc-393900 FITC), Alexa Fluor® 488 (sc-393900 AF488), Alexa Fluor® 546 (sc-393900 AF546), Alexa Fluor® 594 (sc-393900 AF594) or Alexa Fluor® 647 (sc-393900 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-393900 AF680) or Alexa Fluor® 790 (sc-393900 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

α-lactalbumin (H-1) is recommended for detection of α-lactalbumin of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for α-lactalbumin siRNA (h): sc-72407, α-lactalbumin shRNA Plasmid (h): sc-72407-SH and α-lactalbumin shRNA (h) Lentiviral Particles: sc-72407-V.


Molecular Weight of α-lactalbumin: 14 kDa.

Positive Controls: human α-lactalbumin transfected HEK293T whole cell lysate.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



α-lactalbumin (H-1): sc-393900. Western blot analysis of α-lactalbumin expression in non-transfected (A) and human α-lactalbumin transfected (B) HEK293T whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.