

Mo-CD3/CD19/CD45

3 Color FCM Reagent: sc-3967



BACKGROUND

Mouse CD3/CD19/CD45: sc-3967 is a direct immunofluorescence reagent formatted to identify and determine the percentage T lymphocytes and B lymphocytes in erythrocyte-lysed whole blood, based on cell-surface antigen expression. CD3 identifies T lymphocytes and non-covalently associates with either α/β or γ/δ TCR (1). CD45 is a major leukocyte cell surface molecule (2). NK lymphocytes identified as CD3⁻ and CD16⁺ and/or CD56⁺ mediate cytotoxicity against certain tumors and virus infected cells (3). CD19 is present on B lymphocytes during all stages of B cell maturation, but is lost on plasma cells (4). The total population of T lymphocytes and B lymphocytes are used to characterize and monitor some forms of immunodeficiency and autoimmune disease (5,6).

Antigen Expression	Cell Type Identified
CD3+	Mature T Lymphocytes
CD3 ⁻ CD19+	Total B Lymphocytes

STORAGE

Store at 4° C. Do not freeze. Stable for one year from the date of shipment. Protect reagents from prolonged exposure to light.

PRODUCT

Supplied in 1.0 ml of PBS containing 0.1% azide and 0.1% gelatin. Sufficient for 50 tests. This product has been titrated for optimal performance. Recommended use is 20 μ L per test (1×10^6 cells). **For research use only. Not for use in diagnostic procedures.**

INSTRUMENT

Mouse CD3/CD19/CD45: sc-3967 is recommended for use with either a single or dual laser Flow Cytometer fitted with appropriate acquisition and analysis software, such as the FACSCalibur™ Flow Cytometer fitted with CellQuest™ Software by Becton Dickinson.

The flow cytometer must be equipped with a 488 nm laser and must be capable of detecting light scatter (forward and side) and three-color fluorescence with emission detectable in three ranges: 515-545 nm, 562-607 nm and >650 nm, and it must be able to threshold and discriminate using the >650 channel.

Antigen	Clone	Isotype	Label*	Detection Range (nm)
CD3	145-2C11	Armenian Hamster IgG	FITC	515-545
CD19	65	rat IgG _{2a}	PE	562-607
CD45	MB4B4	rat IgG	PE-Cy5	>650

*Fluorescent labels include FITC: Fluorescein isothiocyanate; PE: phycoerythrin; PE-Cy5: phycoerythrin-cyanin 5.

ISOTYPE CONTROL

sc-3967 CON (Armenian Hamster IgG FITC/rat IgG_{2a} PE/rat IgG PE-Cy5) is the isotype matched negative control for this system and is suitable for 50 tests.

REFERENCES

- Exley, M., Terhorst, C., and Wileman, T. 1991. Structure, assembly and intracellular transport of the T cell receptor for antigen. *Semin. Immunol.* 3: 283-297.
- Charbonneau, H., Tonks, N.K., Walsh, K.A., and Fischer, E.H. 1988. The leukocyte common antigen (CD45): a putative receptor-linked protein tyrosine phosphatase. *Proc. Natl. Acad. Sci. USA* 85: 7182-7186.
- Fitzgerald-Bocarsly, P., Herberman, R., Hercend, T., *et al.* 1989. A definition of natural killer cells. In: Ades, E., Lopez, C., eds. *Natural Killer Cells and Host Defense*. Fasel: Karger; 1.
- Dörken, B., Möller, P., Pezzutto, A., Schwartz-Albiez, R., and Moldenhauer, G. B-cell antigens: CD19. In: Knapp, W., Dörken, B., Gilks, W.R. *et al.* eds. *Leucocyte Typing IV: White Cell Differentiation Antigens*. New York, NY: Oxford University Press; 1989: 34-36.
- Foucar, K. and Goeken, J.A. 1982. Clinical Applications of immunologic techniques to the diagnosis of lymphoproliferative and immunodeficiency disorders. *Lab. Med.* 13: 403-413.
- Smolen, J.S., Chused, T.M., Leiserson, W.M., Reeves, J.P., Alling, D., and Steinberg, A.D. 1982. Heterogeneity of immunoregulatory T-cell subsets in systemic lupus erythematosus. Correlation with clinical features. *Am. J. Med.* 2: 783-790.