SANTA CRUZ BIOTECHNOLOGY, INC.

GPI (E-4): sc-398382



BACKGROUND

Glucose-6-phosphate isomerase (GPI) has many other names, including phosphohexose isomerase (PHI), neuroleukin (NLK) and spermantigen-36 (SA-36). GPI is a cytoplasmic homodimer belonging to the GPI family. It is a neurotrophic factor for spinal and sensory neurons and is involved in glycolysis and gluconeogenesis. Defects or mutations in GPI can cause hereditary nonspherocytic hemolytic anemia (HA), hydrops fetalis, immediate neonatal death and neurological impairment.

REFERENCES

- Beutler, E., et al. 1997. Glucosephosphate isomerase (GPI) deficiency mutations associated with hereditary nonspherocytic hemolytic anemia (HNSHA). Blood Cells Mol. Dis. 23: 402-409.
- Kugler, W., et al. 1998. Molecular basis of neurological dysfunction coupled with haemolytic anemia in human glucose-6-phosphate isomerase (GPI) deficiency. Hum. Genet. 103: 450-454.
- Schulz, L.C., et al. 2003. Glucose-6-phosphate isomerase is necessary for embryo implantation in the domestic ferret. Proc. Natl. Acad. Sci. USA 100: 8561-8566.
- Muraki, Y., et al. 2004. Glucose-6-phosphate isomerase variants play a key role in the generation of anti-GPI antibodies: possible mechanism of autoantibody production. Biochem. Biophys. Res. Commun. 323: 518-522.
- Graham Solomons, J.T., et al. 2004. The crystal structure of mouse phosphoglucose isomerase at 1.6A resolution and its complex with glucose-6-phosphate reveals the catalytic mechanism of sugar RING opening. J. Mol. Biol. 342: 847-860.
- Schubert, D., et al. 2004. Immunization with glucose-6-phosphate isomerase induces T cell-dependent peripheral polyarthritis in genetically unaltered mice. J. Immunol. 172: 4503-4509.
- 7. SWISS-PROT/TrEMBL (P06744). World Wide Web URL: http://www.expasy.ch/sprot/sprot-top.html

CHROMOSOMAL LOCATION

Genetic locus: GPI (human) mapping to 19q13.11.

SOURCE

GPI (E-4) is a mouse monoclonal antibody raised against amino acids 1-300 mapping at the N-terminus of GPI of human origin.

PRODUCT

Each vial contains 200 μg lgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GPI (E-4) is recommended for detection of GPI of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPI siRNA (h): sc-43810, GPI shRNA Plasmid (h): sc-43810-SH and GPI shRNA (h) Lentiviral Particles: sc-43810-V.

Molecular Weight (predicted) of GPI: 63 kDa.

Molecular Weight (observed) of GPI: 55 kDa.

Positive Controls: HEK293 whole cell lysate: sc-45136, HeLa whole cell lysate: sc-2200 or RT-4 whole cell lysate: sc-364257.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





GPI (E-4): sc-398382. Western blot analysis of GPI expression in HEK293 (A), HeLa (B), RT-4 (C) and U-251-MG (D) whole cell lysates.

GPI (E-4): sc-398382. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and membrane localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.