

KPI-2 (D-11): sc-398396

BACKGROUND

KPI-2 (kinase/phosphatase/inhibitor 2), also known as BREK, LMR2, cprk, AATYK2 or LMTK2 (lemur tyrosine kinase 2), is a 1,503 amino acid single-pass membrane protein belonging to the protein kinase superfamily and the protein tyrosine kinase family. Mainly expressed in skeletal muscle, and weakly in brain and pancreas, KPI-2 contains N-terminal transmembrane helices and a long C-terminal cytoplasmic tail with serine/threonine/tyrosine kinase activity. KPI-2 may be involved in nerve growth factor (NGF)-Trk A signaling, endosomal membrane trafficking and spermatogenesis. KPI-2 localizes to cytoplasmic membrane vesicles and to perinuclear recycling endosomes. KPI-2 is critical for the transition of endocytosed membrane vesicles from early endosomes to recycling endosomes. The gene encoding KPI-2 is a potential therapeutic target for prostate cancer.

REFERENCES

1. Wang, H., et al. 2002. A novel transmembrane Ser/Thr kinase complexes with protein phosphatase-1 and inhibitor-2. *J. Biol. Chem.* 277: 49605-49612.
2. Kawa, S., et al. 2004. Involvement of BREK, a serine/threonine kinase enriched in brain, in NGF signalling. *Genes Cells* 9: 219-232.
3. Wang, H., et al. 2006. Peptide microarray analysis of substrate specificity of the transmembrane Ser/Thr kinase KPI-2 reveals reactivity with cystic fibrosis transmembrane conductance regulator and phosphorylase. *Mol. Cell. Proteomics* 5: 2124-2130.
4. Kawa, S., et al. 2006. Azoospermia in mice with targeted disruption of the Brek/Lmtk2 (brain-enriched kinase/lemur tyrosine kinase 2) gene. *Proc. Natl. Acad. Sci. USA* 103: 19344-19349.
5. Chibalina, M.V., et al. 2007. Myosin VI and its interacting protein LMTK2 regulate tubule formation and transport to the endocytic recycling compartment. *J. Cell Sci.* 120: 4278-4288.

CHROMOSOMAL LOCATION

Genetic locus: LMTK2 (human) mapping to 7q21.3.

SOURCE

KPI-2 (D-11) is a mouse monoclonal antibody raised against amino acids 15-148 mapping near the N-terminus of KPI-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

KPI-2 (D-11) is available conjugated to agarose (sc-398396 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398396 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398396 PE), fluorescein (sc-398396 FITC), Alexa Fluor® 488 (sc-398396 AF488), Alexa Fluor® 546 (sc-398396 AF546), Alexa Fluor® 594 (sc-398396 AF594) or Alexa Fluor® 647 (sc-398396 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398396 AF680) or Alexa Fluor® 790 (sc-398396 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

KPI-2 (D-11) is recommended for detection of KPI-2 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for KPI-2 siRNA (h): sc-75397, KPI-2 shRNA Plasmid (h): sc-75397-SH and KPI-2 shRNA (h) Lentiviral Particles: sc-75397-V.


Molecular Weight of KPI-2: 250 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, ES-2 cell lysate: sc-24674 or ARPE-19 whole cell lysate: sc-364357.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



KPI-2 (D-11): sc-398396. Western blot analysis of KPI-2 expression in HeLa (A), ARPE-19 (B) and ES-2 (C) whole cell lysates.



KPI-2 (D-11): sc-398396. Immunofluorescence staining of methanol-fixed HeLa cells showing Golgi apparatus and cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA