

# RAP55 (C-5): sc-398552



The Power to Question

## BACKGROUND

RAP55, also known as LSM14A (LSM14 homolog A), is a 463 amino acid member of the LSM14 family that contains a FFD-TFG box domain. The LSM14 domain and the RGG repeats of RAP55 are required for accumulation in P-bodies, and the region containing the FDF motif is responsible for cytoplasmic retention. RAP55, and all other Sm-like proteins, contain the Sm sequence motif, which consists of two regions separated by a linker of variable length that folds as a loop. The Sm-like proteins are thought to form a stable heteromer present in tri-snRNP particles, which are important for pre-mRNA splicing. RAP55 is phosphorylated upon DNA damage, probably by ATM or ATR. Existing as two alternatively spliced isoforms, the RAP55 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken, zebrafish, mosquito, *Drosophila*, *C. elegans*, *S. pombe*, *S. cerevisiae*, *K. lactis*, *E. gossypii*, *M. grisea* and *N. crassa*, and maps to human chromosome 19q13.11.

## CHROMOSOMAL LOCATION

Genetic locus: LSM14A (human) mapping to 19q13.11; Lsm14a (mouse) mapping to 7 B1.

## SOURCE

RAP55 (C-5) is a mouse monoclonal antibody raised against amino acids 103-197 mapping within an internal region of RAP55 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RAP55 (C-5) is available conjugated to agarose (sc-398552 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398552 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398552 PE), fluorescein (sc-398552 FITC), Alexa Fluor® 488 (sc-398552 AF488), Alexa Fluor® 546 (sc-398552 AF546), Alexa Fluor® 594 (sc-398552 AF594) or Alexa Fluor® 647 (sc-398552 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398552 AF680) or Alexa Fluor® 790 (sc-398552 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS

RAP55 (C-5) is recommended for detection of RAP55 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RAP55 siRNA (h): sc-97825, RAP55 siRNA (m): sc-152701, RAP55 shRNA Plasmid (h): sc-97825-SH, RAP55 shRNA Plasmid (m): sc-152701-SH, RAP55 shRNA (h) Lentiviral Particles: sc-97825-V and RAP55 shRNA (m) Lentiviral Particles: sc-152701-V.


Molecular Weight of RAP55: 51 kDa.

Positive Controls: A549 cell lysate: sc-2413, HeLa whole cell lysate: sc-2200 or Hep G2 cell lysate: sc-2227.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



RAP55 (C-5): sc-398552. Western blot analysis of RAP55 expression in Hep G2 (A), A549 (B), HeLa (C), THP-1 (D) and KNRK (E) whole cell lysates. Detection reagent used: m-IgG<sub>1</sub> BP-HRP: sc-525408.



RAP55 (C-5): sc-398552. Western blot analysis of RAP55 expression in A549 (A), KNRK (B) and 3T3-L1 (C) whole cell lysates.

## SELECT PRODUCT CITATIONS

- Mili, D., et al. 2015. Localization and role of RAP55/LSM14 in HeLa cells: a new finding on the mitotic spindle assembly. *Acta Biochim. Pol.* 62: 613-619.
- Samhadaneh, D.M., et al. 2019. Gold nanourchins induce cellular stress, impair proteostasis and damage RNA. *Nanomedicine* 22: 102083.
- Zhang, H., et al. 2024. LSM14B coordinates protein component expression in the P-body and controls oocyte maturation. *J. Genet. Genomics* 51: 48-60.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.