

ATE1 (G-6): sc-398805

BACKGROUND

Arginyl-tRNA-protein transferase (ATE1), also designated arginyltransferase 1, belongs to the R-transferase family of proteins. In order for a protein to be degraded via the ubiquitin pathway, arginylation of the protein is required. ATE1 plays an important role in this process, as it is important for the post-translational conjugation of arginine to the N-terminal aspartate-, glutamate- and possibly cystine-containing substrates. ATE1 is a 518 amino acid protein. Alternative splicing results in two distinct isoforms. ATE1, which is found as a monomer, can localize to the cytoplasm and/or the nucleus.

REFERENCES

1. Kwon, Y.T., et al. 1999. Alternative splicing results in differential expression, activity, and localization of the two forms of arginyl-tRNA-protein transferase, a component of the N-end rule pathway. *Mol. Cell. Biol.* 19: 182-193.
2. Kwon, Y.T., et al. 2002. An essential role of N-terminal arginylation in cardiovascular development. *Science* 297: 96-99.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607103. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Hu, R.G., et al. 2005. The N-end rule pathway as a nitric oxide sensor controlling the levels of multiple regulators. *Nature* 437: 981-986.
5. Lee, M.J., et al. 2005. RGS4 and RGS5 are *in vivo* substrates of the N-end rule pathway. *Proc. Natl. Acad. Sci. USA* 102: 15030-15035.

CHROMOSOMAL LOCATION

Genetic locus: ATE1 (human) mapping to 10q26.13; Ate1 (mouse) mapping to 7 F3.

SOURCE

ATE1 (G-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 244-269 within an internal region of ATE1 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ATE1 (G-6) is available conjugated to agarose (sc-398805 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-398805 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-398805 PE), fluorescein (sc-398805 FITC), Alexa Fluor® 488 (sc-398805 AF488), Alexa Fluor® 546 (sc-398805 AF546), Alexa Fluor® 594 (sc-398805 AF594) or Alexa Fluor® 647 (sc-398805 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-398805 AF680) or Alexa Fluor® 790 (sc-398805 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-398805 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

ATE1 (G-6) is recommended for detection of ATE1 isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ATE1 siRNA (h): sc-60220, ATE1 siRNA (m): sc-60221, ATE1 shRNA Plasmid (h): sc-60220-SH, ATE1 shRNA Plasmid (m): sc-60221-SH, ATE1 shRNA (h) Lentiviral Particles: sc-60220-V and ATE1 shRNA (m) Lentiviral Particles: sc-60221-V.


Molecular Weight of ATE1: 59 kDa.

Positive Controls: U-251-MG whole cell lysate: sc-364176 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



ATE1 (G-6): sc-398805. Western blot analysis of ATE1 expression in Hep G2 (A) and U-251-MG (B) whole cell lysates.



ATE1 (G-6): sc-398805. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

1. Kasu, Y.A.T., et al. 2018. The N-termini of TAR DNA-binding protein-43 (TDP43) C-terminal fragments influence degradation, aggregation propensity and morphology. *Mol. Cell. Biol.* 38: e00243-18.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.