

Syntaxin 7 siRNA (h): sc-41334

BACKGROUND

Correct vesicular transport is essential to the survival of eukaryotic cells. This process is determined by specific pairing of vesicle-associated SNAREs (v-SNAREs) with those on the target membrane (t-SNAREs). This complex then recruits soluble NSF attachment proteins (SNAPs) and N-ethylmaleimide-sensitive factor (NSF) to form the highly stable SNAP receptor (SNARE) complex. The formation of a SNARE complex pulls the vesicle and target membrane together and may provide the energy to drive fusion of the lipid bilayers. Syntaxins, a family of proteins involved in the fusion of synaptic vesicles with the plasma membrane, display broad tissue distribution and contain carboxy-terminal hydrophobic domains that direct themselves to their respective intracellular compartments. Syntaxin 7 binds α -SNAP *in vitro* and forms a complex with Syntaxin 8, vti1b and VAMP-8 that functions in the fusion of late endosomes. *In vitro*, the abundant expression of Syntaxin 7 in B16 melanoma cells increases as they undergo melanogenesis. A SNARE complex between Syntaxin 7 and VAMP7 or VAMP8 may regulate the fusion events that eventually lead to melanogenesis.

REFERENCES

1. Elferink, L.A., et al. 1993. A role for Synaptotagmin (p65) in regulated exocytosis. *Cell* 72: 153-159.
2. Bennett, M.K., et al. 1993. The Syntaxin family of vesicular transport receptors. *Cell* 74: 863-873.
3. Yamaguchi, K., et al. 1994. Exocytosis relating proteins in the nervous system. *Neurosci. Res.* 20: 289-292.
4. Hayashi, T., et al. 1994. Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on assembly. *EMBO J.* 13: 5051-5061.
5. Edelmann, L., et al. 1995. Synaptobrevin binding to synaptophysin: a potential mechanism for controlling the exocytosis fusion machine. *EMBO J.* 14: 224-231.
6. Lin, R.C. and Scheller, R.H. 1997. Structural organization of the synaptic exocytosis core complex. *Neuron* 19: 1087-1094.
7. Barnard, R.J., et al. 1997. Stimulation of NSF ATPase activity by α -SNAP is required for SNARE complex disassembly and exocytosis. *J. Cell Biol.* 139: 875-883.

CHROMOSOMAL LOCATION

Genetic locus: STX7 (human) mapping to 6q23.2.

PRODUCT

Syntaxin 7 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Syntaxin 7 shRNA Plasmid (h): sc-41334-SH and Syntaxin 7 shRNA (h) Lentiviral Particles: sc-41334-V as alternate gene silencing products.

For independent verification of Syntaxin 7 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41334A, sc-41334B and sc-41334C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Syntaxin 7 siRNA (h) is recommended for the inhibition of Syntaxin 7 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Syntaxin 7 (A-1): sc-514017 is recommended as a control antibody for monitoring of Syntaxin 7 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Syntaxin 7 gene expression knockdown using RT-PCR Primer: Syntaxin 7 (h)-PR: sc-41334-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.