

GABA T-2 siRNA (m): sc-41961

BACKGROUND

GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (γ -aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl^- conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic ($GABA_A$) and metabotropic ($GABA_B$) receptors as well as a third class of receptors called $GABA_C$. Both $GABA_A$ and $GABA_C$ are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the $GABA_A$ receptor family include $GABA_A R\alpha 1-6$, $GABA_A R\beta 1-3$, $GABA_A R\gamma 1-3$, $GABA_A R\delta$, $GABA_A R\epsilon$, $GABA_A R\rho 1$ and $GABA_A R\rho 2$. The $GABA_B$ family is composed of $GABA_B R1\alpha$ and $GABA_B R1\beta$. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

REFERENCES

- Nelson, H., et al. 1990. Cloning of the human brain GABA transporter. *FEBS Lett.* 269: 181-184.
- Cherubini, E., et al. 1991. GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci.* 14: 515-519.
- Borden, L.A., et al. 1992. Molecular heterogeneity of the γ -aminobutyric acid (GABA) transport system. Cloning of two novel high affinity GABA transporters from rat brain. *J. Biol. Chem.* 267: 21098-21104.
- Dirx, R., Jr., et al. 1995. Targeting of the 67-kDa isoform of glutamic acid decarboxylase to intracellular organelles is mediated by its interaction with the NH_2 -terminal region of the 65-kDa isoform of glutamic acid decarboxylase. *J. Biol. Chem.* 270: 2241-2246.
- Lukasiewicz, P.D. 1996. $GABA_C$ receptors in the vertebrate retina. *Mol. Neurobiol.* 12: 181-194.
- Kaupmann, K., et al. 1997. Expression cloning of $GABA_B$ receptors uncovers similarity to metabotropic glutamate receptors. *Nature* 386: 239-246.
- Korpi, E.R., et al. 1997. $GABA_A$ -receptor subtypes: clinical efficiency and selectivity of benzodiazepine site ligands. *Ann. Med.* 29: 275-282.

CHROMOSOMAL LOCATION

Genetic locus: Slc6a13 (mouse) mapping to 6 F1.

PRODUCT

GABA T-2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GABA T-2 shRNA Plasmid (m): sc-41961-SH and GABA T-2 shRNA (m) Lentiviral Particles: sc-41961-V as alternate gene silencing products.

For independent verification of GABA T-2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41961A, sc-41961B and sc-41961C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at $-20^\circ C$ with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at $-20^\circ C$, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GABA T-2 siRNA (m) is recommended for the inhibition of GABA T-2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GABA T-2 gene expression knockdown using RT-PCR Primer: GABA T-2 (m)-PR: sc-41961-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ C$ and the extension temperature should be $68-72^\circ C$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.