

Skp1 p19 (1-162): sc-4230 WB

BACKGROUND

The critical role that the family of regulatory proteins known as cyclins plays in eukaryotic cell cycle regulation is well established. The best characterized cyclin complex is the mitotic cyclin B/Cdc2 p34 kinase, the active component of M phase promoting factor. Cyclin A accumulates prior to cyclin B in the cell cycle, appears to be involved in control of S phase and has been shown to associate with cyclin dependent kinase-2 (Cdk2). In addition, cyclin A has been implicated in cell transformation and is found in complexes with E1A, transcription factors DP-1 and E2F and retinoblastoma protein p110. Two cyclin A-Cdk2 complex binding proteins, Skp1 p19 and Skp2 p45, have been described. Although the Skps (S phase kinase-associated proteins) associate with the active cyclin A-Cdk2 complex, they do not exhibit any regulatory effects on the complex. Abolition of Skp2 p45 function by either microinjection of anti-p45 antibodies or addition of antisense oligonucleotides prevents entry into S phase of both normal and transformed cells.

REFERENCES

1. Draetta, G., Lucas, F., Westendorf, J., Brizuela, L., Ruderman, J., and Beach D. 1989. Cdc2 protein kinase is complexed with both cyclin A and B: evidence for proteolytic inactivation of MPF. *Cell* 56: 829-838.
2. Giordano, A., Whyte, P., Harlow, E., Franza, B.R. Jr., Beach, D., and Draetta, G. 1989. A 60 kd Cdc2-associated polypeptide complexes with the E1A proteins in adenovirus-infected cells. *Cell* 58: 981-990.
3. Pines, J. and Hunter T. 1990. Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* 346: 760-763.
4. Wang, J., Chenivasse, X., Henglein, B., and Berchot, C. 1990. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 343: 555-557.
5. Bandara, L.R., Adamczewski, J.P., Hunt, T., and La Thangue, N.B. 1991. Cyclin A and the retinoblastoma gene product complex with a common transcription factor. *Nature* 352: 249-251.
6. Williams, R.T., Carbonaro-Hall, D., and Hall, F.L. 1992. Co-purification of p34Cdc2/p58 cyclin A proline-directed protein kinase and the retinoblastoma tumor susceptibility gene product: interaction of an oncogenic serine/threonine protein kinase with a tumor-suppressor protein. *Oncogene* 7: 423-432.
7. Lees, E., Faha, B., Dulic, V., Reed, S.I., and Harlow, E. 1992. Cyclin E/Cdk2 and cyclin A/Cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* 6: 1874-1885.
8. Faha, B., Ewen, M., Tsai, L.H., Livingston, D., and Harlow, E. 1992. Interaction between human cyclin A and adenovirus E1A-associated p107 protein. *Science* 255: 87-90.
9. Zhang, H., Kobayashi, R., Galaktionov, K., and Beach, D. 1995. p19Skp1 and p45Skp2 are essential elements of the cyclin A-Cdk2 S phase kinase. *Cell* 82: 915-925.

STORAGE

Store at -20° C; stable for one year from the date of shipment.

SOURCE

Skp1 p19 (1-162) is expressed in *E. coli* as a 46 kDa tagged fusion protein of human origin corresponding to amino acids 1-162 representing full length Skp1 p19.

PRODUCT

Skp1 p19 (1-162) is purified from bacterial lysates by glutathione agarose affinity chromatography; supplied as 10 µg in 0.1 ml SDS-PAGE loading buffer.

APPLICATIONS

Skp1 p19 (1-162) is suitable as a Western blotting control for sc-1568, sc-1569 and sc-7163.

RESEARCH USE

For research use only, not for use in diagnostic procedures.