

AQP2 siRNA (m): sc-42364

BACKGROUND

Aquaporins (AQPs) are a large family of integral membrane water transport channel proteins that facilitate the transport of water through the cell membrane. This function is conserved in animals, plants and bacteria. Many isoforms of aquaporin have been identified in mammals, designated AQP0 through AQP10. Aquaporins are widely distributed and it is not uncommon for more than one type of AQP to be present in the same cell. Although most aquaporins are only permeable to water, AQP3, AQP7, AQP9 and one of the two AQP10 transcripts are also permeable to urea and glycerol. AQP2 is the only water channel that is activated by Vasopressin to enhance water reabsorption in the kidney collecting duct. Aquaporins are involved in renal water absorption, generation of pulmonary secretions, lacrimation, and the secretion and reabsorption of cerebrospinal fluid and aqueous humor.

REFERENCES

1. Preston, G.M., et al. 1991. Isolation of the cDNA for erythrocyte integral membrane protein of 28 kilodaltons: member of an ancient channel family. *Proc. Natl. Acad. Sci. USA* 88: 11110-11114.
2. Sasaki, S., et al. 1993. Cloning, expression and chromosomal mapping of human collecting duct water channel (hWCH-CD). *J. Am. Soc. Nephrol.* 4: 858.
3. Deen, P.M., et al. 1994. Requirement of human renal water channel aquaporin 2 for Vasopressin-dependent concentration of urine. *Science* 264: 92-95.
4. Ishibashi, K., et al. 1995. Structure and chromosomal localization of a human water channel (AQP3) gene. *Genomics* 27: 352-354.
5. Yang, B., et al. 1995. cDNA cloning, gene organization, and chromosomal localization of a human mercurial insensitive water channel. Evidence for distinct transcriptional units. *J. Biol. Chem.* 270: 22907-22913.
6. Lee, M.D., et al. 1996. The human aquaporin-5 gene. Molecular characterization and chromosomal localization. *J. Biol. Chem.* 271: 8599-8604.
7. Valenti, G., et al. 1996. Expression and functional analysis of water channels in a stably AQP2-transfected human collecting duct cell line. *J. Biol. Chem.* 271: 24365-24370.
8. Echevarria, M., et al. 1998. Aquaporins. *J. Physiol. Biochem.* 54: 107-118.
9. Beitz, E., et al. 1999. The mammalian aquaporin water channel family: a promising new drug target. *Curr. Med. Chem.* 6: 457-467.

CHROMOSOMAL LOCATION

Genetic locus: *Aqp2* (mouse) mapping to 15 F1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

AQP2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AQP2 shRNA Plasmid (m): sc-42364-SH and AQP2 shRNA (m) Lentiviral Particles: sc-42364-V as alternate gene silencing products.

For independent verification of AQP2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42364A, sc-42364B and sc-42364C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AQP2 siRNA (m) is recommended for the inhibition of AQP2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

AQP2 (E13-21): sc-47710 is recommended as a control antibody for monitoring of AQP2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AQP2 gene expression knockdown using RT-PCR Primer: AQP2 (m)-PR: sc-42364-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.