

KIR6.2 siRNA (h): sc-42628

BACKGROUND

ATP-sensitive K⁺ channels play important roles in many cellular functions by coupling cell metabolism to electrical activity. KIR6.1 and KIR6.2 are members of the KIR (inwardly rectifying potassium channel) family of potassium channels. Inward rectifying K⁺ channels possess a greater tendency to allow potassium to flow into the cell rather than out of it. These channels comprise two subunits: a KIR6.0 subfamily component and a SUR component, which is a member of the ATP-binding cassette protein superfamily. Mutations in the gene coding for these channels are a cause of an autosomal recessive disorder characterized by unregulated Insulin secretion. The amino-terminal and carboxyl-terminal domains of KIR channel subunits are both intracellular, and the two intracellular domains of KIR6.2 physically interact with each other.

REFERENCES

1. Inagaki, N., et al. 1995. Reconstitution of IKATP: an inward rectifier subunit plus the sulfonylurea receptor. *Science* 270: 1166-1170.
2. Isomoto, S., et al. 1997. Inwardly rectifying potassium channels: their molecular heterogeneity and function. *Jpn. J. Physiol.* 47: 11-39.
3. Inagaki, N., et al. 1998. ATP-sensitive potassium channels: structures, functions, and pathophysiology. *Jpn. J. Physiol.* 48: 397-412.
4. Tucker, S.J., et al. 1999. Mapping of the physical interaction between the intracellular domains of an inwardly rectifying potassium channel, KIR6.2. *J. Biol. Chem.* 274: 33393-33397.
5. Meissner, T., et al. 1999. Congenital hyperinsulinism: molecular basis of a heterogeneous disease. *Hum. Mutat.* 13: 351-361.
6. Miki, T., et al. 1999. The structure and function of the ATP-sensitive K⁺ channel in Insulin-secreting pancreatic β cells. *J. Mol. Endocrinol.* 22: 113-123.
7. Seino, S. 1999. ATP-sensitive potassium channels: a model of hetero-multimeric potassium channel/receptor assemblies. *Annu. Rev. Physiol.* 61: 337-362.

CHROMOSOMAL LOCATION

Genetic locus: KCNJ11 (human) mapping to 11p15.1.

PRODUCT

KIR6.2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIR6.2 shRNA Plasmid (h): sc-42628-SH and KIR6.2 shRNA (h) Lentiviral Particles: sc-42628-V as alternate gene silencing products.

For independent verification of KIR6.2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42628A, sc-42628B and sc-42628C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KIR6.2 siRNA (h) is recommended for the inhibition of KIR6.2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

KIR6.2 (B-9): sc-390104 is recommended as a control antibody for monitoring of KIR6.2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG λ BP-HRP: sc-516132 or m-IgG λ BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG λ BP-FITC: sc-516185 or m-IgG λ BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIR6.2 gene expression knockdown using RT-PCR Primer: KIR6.2 (h)-PR: sc-42628-PR (20 μ l, 546 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.