

CD8- β (22-170): sc-4295 WB

BACKGROUND

The T cell receptor (TCR) is a heterodimer composed of either α and β or γ and δ chains. CD3 chains and the CD4 or CD8 co-receptors are also required for efficient signal transduction through the TCR. The TCR is expressed on T helper and T cytotoxic cells that can be distinguished by their expression of CD4 and CD8. T helper cells express CD4 proteins and T cytotoxic cells display CD8. CD8, also designated Leu 2 or T8, is a 32 kDa cell surface glycoprotein. It is a two chain complex (α - α or α - β) receptor that binds class I MHC molecules presented by the antigen-presenting cell (APC). A primary function of CD8 is to facilitate antigen recognition by the TCR and to strengthen the avidity of the TCR-antigen interactions. An additional role for CD8-expressing T cells may be to maintain low levels of HIV expression.

REFERENCES

1. Nakayama, K., Tokito, S., Okumura, K. and Nakauchi, H. 1989. Structure and expression of the gene encoding CD8 α chain (Leu-2/T8). Immunogenetics 30: 393-397.
2. Fleury, S.G., Croteau, G. and Sekaly, R.P. 1991. CD4 and CD8 recognition of class II and class I molecules of the major histocompatibility complex. Sem. Immunol. 3: 177-185.
3. Allison, J.P. and Havran, W.L. 1991. The immunobiology of T cells with invariant $\gamma\delta$ antigen regions. Annu. Rev. Immunol. 9: 679-705.
4. Zuniga-Pflucker, J.C., Jones, L.A., Chin, L.T. and Kruisbeek, A.M. 1991. CD4 and CD8 act as coreceptors during thymic selection of the T cell repertoire. Sem. Immunol. 3: 167-175.
5. Janeway, C.A., Jr. 1992. The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. Annu. Rev. Immunol. 10: 645-674.
6. Buseyne, F. and Riviere, Y. 1993. HIV-specific CD8⁺ T-cell immune responses and viral replication. AIDS 2 Suppl.: S81-S85.
7. Julius, M., Maroun, C.R. and Haughn, L. 1993. Distinct roles for CD4 and CD8 as co-receptors in antigen receptor signalling. Immunol. Today 14: 177-183.
8. Ehrlich, E.W., Devaux, B., Rock, E.P., Jorgenson, J.L., Davis, M.N. and Chien, Y.H. 1993. T cell receptor interaction with peptide/major histocompatibility complex (MHC) and superantigen MHC ligands is dominated by antigen. J. Exp. Med. 178: 713-722.
9. Luescher, I.F., Vivier, E., Layer, A., Mahiou, J., Godeau, F., Malissen, B. and Romero, P. 1995. CD8 modulation of T-cell antigen receptor-ligand interactions on living cytotoxic T lymphocytes. Nature 373: 353-356.

SOURCE

CD8- β (22-170) is expressed in *E. coli* as a 43 kDa tagged fusion protein corresponding to amino acids 22-170 of CD8- β of human origin.

PRODUCT

CD8- β (22-170) is purified from bacterial lysates (>98%) by glutathione agarose affinity chromatography; supplied as 10 μ g in 0.1 ml SDS-PAGE loading buffer.

APPLICATIONS

CD8-b (22-170) is suitable as a Western blotting control for sc-9147.

STORAGE

Store at -20° C; stable for one year from the date of shipment.

RESEARCH USE

For research use only, not for use in diagnostic procedures.