



Sox-15 (121-233): sc-4447 WB

BACKGROUND

Sox-20 (SRY (sex determining region Y)-box 20, SRY-box 15, SOX26, SOX27) encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Sox-20 may act as a transcriptional regulator after forming a protein complex with other proteins. Sox-20 is widely expressed in fetal and adult tissues with highest level found in fetal spinal cord, adult brain and adult testis. Sox family transcription factors influence cell differentiation, development, and sex determination. Sox-20 contains a unique DNA binding domain known as the high mobility group (HMG) box that is related to that of the testis determining gene SRY. The highly complex group of Sox genes cluster at a minimum of 40 different loci that rapidly diverged in various animal lineages. Several Sox genes have been identified, and members of this family have been shown to be conserved during evolution and to play key roles during animal development.

REFERENCES

1. Laudet, V., Stehelin, D., and Clevers, H. 1993. Ancestry and diversity of the HMG box superfamily. *Nucleic Acids Res.* 21: 2493-2501.
2. Meyer, J., Wirth, J., Held, M., Schempp, W., and Scherer, G. 1996. Sox20, a new member of the SOX gene family, is located on chromosome 17p13. *Cytogenet. Cell Genet.* 72: 246-249.
3. Kuhlbrodt, K., Herbarth, B., Sock, E., Hermans-Borgmeyer, I., and Wegner, M. 1998. Sox10, a novel transcriptional modulator in glial cells. *J. Neurosci.* 18: 237-250.
4. Critcher, R., Stitson, R.N., Wade-Martins, R., Easty, D.J., and Farr, C.J. 1998. Assignment of Sox4 to mouse chromosome 13 bands A3-A5 by fluorescence *in situ* hybridization; refinement of the human Sox4 location to 6p22.3 and of Sox20 to chromosome 17p12.3. *Cytogenet. Cell Genet.* 81: 294-295
5. Arsic, N., Rajic, T., Stanojic, S., Goodfellow, P.N., and Stevanovic, M. 1998. Characterisation and mapping of the human Sox14 gene. *Cytogenet. Cell Genet.* 83: 139-146.
6. Vujic, M., Rajic, T., Goodfellow, P.N., and Stevanovic, M. 1998. cDNA characterization and high resolution mapping of the human Sox20 gene. *Mamm Genome* 9: 1059-1061.
7. Hiraoka, Y., Ogawa, M., Sakai, Y., Taniguchi, K., Fujii, T., Umezawa, A., Hata, J., and Aiso, S. 1998. Isolation and expression of a human SRY-related cDNA hSox20. *Biochim. Biophys. Acta.* 1396: 132-137.
8. Osaki, E., Nishina, Y., Inazawa, J., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Ohsugi, M., Tezuka, T., Yoshida, M., and Semba, K. 1999. Identification of a novel Sry-related gene and its germ cell-specific expression. *Nucleic Acids Res.* 27: 2503-2510.
9. Sasai, Y. 2001. Roles of Sox factors in neural determination: conserved signaling in evolution? *Int. J. Dev. Biol.* 45: 321-326.

SOURCE

Sox-15 (121-233) is expressed in *E. coli* as a 40 kDa tagged fusion protein corresponding to amino acids 121-233 of Sox-15 of human origin.

PRODUCT

Sox-15 (121-233) is purified from bacterial lysates (>98%) by glutathione agarose affinity chromatography; supplied as 10 µg in 0.1 ml SDS-PAGE loading buffer.

APPLICATIONS

SOX-15 (121-233) is suitable as a Western blotting control for sc-20101.

STORAGE

Store at -20° C; stable for one year from the date of shipment.

RESEARCH USE

For research use only, not for use in diagnostic procedures.