

donkey anti-rabbit IgG-PerCP: sc-45095

BACKGROUND

Santa Cruz Biotechnology's secondary antibodies are available conjugated to either an enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, immunostaining, flow cytometry and ELISA. Secondary antibodies are commonly affinity purified against immobilized whole IgG or against antibody fragments. Santa Cruz Biotechnology offers an extensive selection of secondary antibodies optimized for immunohistochemistry and flow cytometry, and are labeled with either biotin, FITC (fluorescein isothiocyanate), Texas Red®, TRITC (tetramethyl rhodamine iso-thiocyanate), PE (phycoerythrin), PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex with cyanin-5.5). Immunohistochemistry and flow cytometry secondary antibodies are specific for commonly used primary antibody species, including goat, rabbit, mouse and rat.

SOURCE

donkey anti-rabbit IgG-PerCP is a pre-adsorbed, affinity purified secondary antibody raised in donkey against rabbit IgG and conjugated to PerCP (peridinin chlorophyll protein complex).

PRODUCT

Each vial contains 200 µg donkey IgG (pre-adsorbed with mouse and human IgG) in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

APPLICATIONS

donkey anti-rabbit IgG-PerCP is recommended for detection of rabbit IgG by immunofluorescence staining (starting dilution: 1:100, dilution range: 1:100-1:400), immunohistochemical staining (starting dilution: 1:100, dilution range: 1:100-1:400) and flow cytometry (0.5-1 µg per 1 x 10⁶ cells).

RECOMMENDED SUPPORT PRODUCTS

A. TISSUE CULTURE CELLS

- CrystalCruz™ Cover Glasses, 22 x 50 mm: sc-24975
- CrystalCruz™ Micro Slides 75 x 25 mm; frosted sides: sc-24976
- PBS, powder: sc-24947
- Formaldehyde: sc-203049
- Hydrogen Peroxide: sc-203336

B. FROZEN TISSUE SECTIONS

- Organo/Limonene Mount: sc-45087
- UltraCruz™ Mounting Medium: sc-24941
- ImmunoHistoMount: sc-45086
- Immuno In Situ Mount: sc-45088

C. FORMALIN-FIXED, PARAFFIN-EMBEDDED TISSUE SECTIONS


- Paraffin: sc-286633
- Xylenes: sc-237422
- Hematoxylin: sc-24973

Texas Red® is a registered trademark of Molecular Probes (6/02).


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



mouse anti-rabbit IgG-PerCP: sc-45098. Indirect, intracellular FCM analysis of fixed and permeabilized Jurkat cells stained with HEB (A-20), followed by PerCP-conjugated mouse anti-rabbit IgG: sc-45098. Black line histogram represents the isotype control, normal rabbit IgG: sc-3888. Antibody tested: HEB (A-20): sc-357.



donkey anti-rabbit IgG-PerCP: sc-45095. Indirect FCM analysis of BJAB cells stained with CD10 (H-321), followed by PerCP-conjugated donkey anti-rabbit IgG: sc-45095. Black line histogram represents the isotype control, normal rabbit IgG: sc-3888. Antibody tested: CD10 (H-321): sc-9149.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.