

FGF-6 (hBA-168): sc-4576

BACKGROUND

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also referred to as basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int-2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10. Members of the FGF family share 30-55% amino acid sequence identity, similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

REFERENCES

- Moore, R., Casey, G., Brookes, S., Dixon, M., Peters, G., and Dickson, C. 1986. Sequence, topography and protein coding potential of mouse Int-2: a putative oncogene activated by mouse mammary tumor virus. *EMBO J.* 5: 919-924.
- Delli Bovi, P., Curatola, A.M., Kern, F.G., Greco, A., Ittmann, M., and Basilico, C. 1987. An oncogene isolated by transfection of Kaposi's sarcoma DNA encodes a growth factor that is a member of the FGF family. *Cell* 50: 729-737.
- Zhan, X., Bates, B., Hu, X., and Goldfarb, M. 1988. The human FGF-5 oncogene encodes a novel protein related to fibroblast growth factors. *Mol. Cell. Biol.* 8: 3487-3495.
- Marics, I., Adelaide, J., Raybaud, F., Mattei, M., Coulier, F., Planche, J., Lapeyriere, O., and Birnbaum, D. 1989. Characterization of the HST-related FGF.6 gene, a new member of the fibroblast growth factor gene family. *Oncogene* 4: 335-340.
- Rifkin, D.B. and Moscatelli, D. 1989. Recent developments in the cell biology of fibroblast growth factor. *J. Cell Biol.* 109: 1-6.
- Dionne, C.A., Crumley, G., Bellot, F., Kaplow, J.M., Searfoss, G., Ruta, M., Burgess, W.H., Jaye, M., and Schlessinger, J. 1990. Cloning and expression of two distinct high-affinity receptors cross-reacting with acidic and basic fibroblast growth factors. *EMBO J.* 9: 2685-2692.
- Tanaka, A., Miyamoto, K., Minamino, N., Takeda, M., Sato, B., Matsuo, H., and Matsumoto, K. 1992. Cloning and characterization of an androgen-induced growth factor essential for the androgen-dependent growth of mouse mammary carcinoma cells. *Proc. Natl. Acad. Sci. USA* 89: 8928-8932.
- Miyamoto, M., Naruo, K., Seko, C., Matsumoto, S., Kondo, T., and Kurokawa, T. 1993. Molecular cloning of a novel cytokine cDNA encoding the ninth member of the fibroblast growth factor family, which has a unique secretion property. *Mol. Cell. Biol.* 13: 4251-4259.
- Beer, H.D., Florence, C., Dammeier, J., McGuire, L., Werner, S., and Duan, D.R. 1997. Mouse fibroblast growth factor 10: cDNA cloning, protein characterization, and regulation of mRNA expression. *Oncogene* 15: 2211-2218.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

FGF-6 (hBA-168) is produced in *E. coli* as 18.7 kDa biologically active protein corresponding to 168 amino acids of FGF-6 of human origin.

PRODUCT

FGF-6 (hBA-168) is purified from bacterial lysates (>98%); supplied as 25 µg purified protein.

BIOLOGICAL ACTIVITY

FGF-6 (hBA-168) is biologically active as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors.

Expected ED₅₀: <0.5 ng/ml.

Specific Activity: Greater than 2 x 10⁶ units/mg

RECONSTITUTION

In order to avoid freeze/thaw damaging of the active protein, dilute protein when first used to desired working concentration. Either a sterile filtered standard buffer (such as 50mM TRIS or 1X PBS) or water can be used for the dilution. Store any thawed aliquot in refrigeration at 2° C to 8° C for up to four weeks, and any frozen aliquot at -20° C to -80° C for up to one year. It is recommended that frozen aliquots be given an amount of standard cryoprotective (such as Ethylene Glycol or Glycerol 5-20% v/v), and refrigerated samples be given an amount of carrier protein (such as heat inactivated FBS or BSA to 0.1% v/v) or non-ionic detergent (such as Triton X-100 or Tween 20 to 0.005% v/v), to aid stability during storage.

STORAGE

Store desiccated at -20° C; stable for one year from the date of shipment.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.