

# HIF PHD1 (A-16): sc-46024

## BACKGROUND

Prolyl hydroxylase domain proteins PHD1, PHD2 and PHD3 (also designated EGLN2, EGLN1, and EGLN3 respectively) can hydroxylate HIF- $\alpha$  subunits. Hypoxia-inducible factor (HIF) is a transcriptional regulator important in several aspects of oxygen homeostasis. The prolyl hydroxylases catalyze the posttranslational formation of 4-hydroxyproline in HIF- $\alpha$  proteins. PHD1, which is widely expressed but in highest levels in testis, functions as a cellular oxygen sensor and is important in cell growth regulation. PHD1, which can localize to the nucleus or the cytoplasm, is also detected in hormone responsive tissues, such as normal and cancerous mammary, ovarian and prostate epithelium. PHD1 is encoded by EGLN2 which maps to chromosome 19q13.2. PHD2 is regarded as the main cellular oxygen sensor as RNA interference against PHD2, but not PHD1 or PHD3, is enough to stabilize HIF-1 $\alpha$  in normoxia. PHD2, a direct HIF target gene, is expressed mainly in skeletal muscle, heart, kidney and brain. PHD3 may play a role in the regulation of cell growth in muscle cells and in apoptosis in neuronal tissue. PHD3 is widely expressed although the highest levels can be detected in placenta and heart.

## CHROMOSOMAL LOCATION

Genetic locus: EGLN2 (human) mapping to 19q13.2; EglN2 (mouse) mapping to 7 A3.

## SOURCE

HIF PHD1 (A-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HIF PHD1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46024 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

HIF PHD1 (A-16) is recommended for detection of HIF PHD1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HIF PHD1 (A-16) is also recommended for detection of HIF PHD1 in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for HIF PHD1 siRNA (h): sc-45616, HIF PHD1 siRNA (m): sc-45617, HIF PHD1 shRNA Plasmid (h): sc-45616-SH, HIF PHD1 shRNA Plasmid (m): sc-45617-SH, HIF PHD1 shRNA (h) Lentiviral Particles: sc-45616-V and HIF PHD1 shRNA (m) Lentiviral Particles: sc-45617-V.

Positive Controls: rat heart extract: sc-2393, mouse heart extract: sc-2254 or human heart extract: sc-363763.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



HIF PHD1 (A-16): sc-46024. Western blot analysis of HIF PHD1 expression in mouse heart (A) and human heart (B) tissue extracts.



HIF PHD1 (A-16): sc-46024. Western blot analysis of HIF PHD1 expression in rat heart tissue extract.

## SELECT PRODUCT CITATIONS

- Lin, H.H., et al. 2011. Andrographolide down-regulates hypoxia-inducible factor-1 $\alpha$  in human non-small cell lung cancer A549 cells. *Toxicol. Appl. Pharmacol.* 250: 336-345.
- Song, X., et al. 2013. Wogonin inhibits tumor angiogenesis via degradation of HIF-1 $\alpha$  protein. *Toxicol. Appl. Pharmacol.* 271: 144-155.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **HIF PHD1 (2G3): sc-293220**, our highly recommended monoclonal alternative to HIF PHD1 (A-16).