

Dok-1 (P-19): sc-46096

BACKGROUND

Dok-1 associates with the Ras GTPase-activating protein (Ras GAP) upon tyrosine phosphorylation. Evidence suggests that Dok-1 (also designated p62dok) is a substrate of the constitutive tyrosine kinase activity of p210 Bcr-Abl, a fusion protein caused by the t(9;22) translocation and associated with chronic myelogenous leukemia. Dok-1, as well as the tyrosine kinase substrates IRS-1 and Cas, are members of a class of "docking" proteins which contain multiple tyrosine residues and putative SH2 binding sites. Dok-1 is suspected to be the substrate phosphorylated in response to stimulation by a number of growth factors, including PDGF, VEGF, Insulin and IGF. Dok-2 (also designated p56dok) has also been identified as a potential mediator of the effects of p210 Bcr-Abl.

REFERENCES

1. Wisniewski, D., et al. 1994. A 62 kDa tyrosine phosphoprotein constitutively present in primary chronic phase chronic myelogenous leukemia enriched lineage negative blast populations. *Leukemia* 8: 688-693.
2. Myers, M.G., et al. 1994. The IRS-1 signaling system. *Trends Biochem. Sci.* 19: 289-293.
3. Guo, D., et al. 1995. Vascular endothelial cell growth factor promotes tyrosine phosphorylation of mediators of signal transduction that contain SH2 domains. Association with endothelial cell proliferation. *J. Biol. Chem.* 270: 6729-6733.
4. Mayer, B.J., et al. 1995. Evidence that SH2 domains promote processive phosphorylation by protein-tyrosine kinases. *Curr. Biol.* 5: 296-305.
5. Holgado, M.M., et al. 1996. A Grb2-associated docking protein in EGF and Insulin receptor signalling. *Nature* 379: 560-564.
6. Carpino, N., et al. 1997. p62dok: a constitutively tyrosine-phosphorylated, GAP-associated protein in chronic myelogenous leukemia progenitor cells. *Cell* 88: 197-204.
7. Yamanashi, Y., et al. 1997. Identification of the Abl- and Ras GAP-associated 62 kDa protein as a docking protein, Dok. *Cell* 88: 205-211.

CHROMOSOMAL LOCATION

Genetic locus: DOK1 (human) mapping to 2p13.1.

SOURCE

Dok-1 (P-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Dok-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46096 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Dok-1 (P-19) is recommended for detection of Dok-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Dok-1 siRNA (h): sc-35210, Dok-1 shRNA Plasmid (h): sc-35210-SH and Dok-1 shRNA (h) Lentiviral Particles: sc-35210-V.


Molecular Weight of Dok-1: 62 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HEL 92.1.7 cell lysate: sc-2270 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Dok-1 (P-19): sc-46096. Western blot analysis of Dok-1 expression in Jurkat (A), K-562 (B) and HEL 92.1.7 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Dok-1 (A-3): sc-6929** or **Dok-1 (45): sc-135888**, our highly recommended monoclonal alternatives to Dok-1 (P-19).