

HSP 20 (L-13): sc-46140

BACKGROUND

The heat shock proteins (HSPs) comprise a group of highly conserved, abundantly expressed proteins with diverse functions, including the assembly and sequestering of multiprotein complexes, transportation of nascent polypeptide chains across cellular membranes and regulation of protein folding. Heat shock proteins (also known as molecular chaperones) fall into six general families: HSP 90, HSP 70, HSP 60, the low molecular weight HSPs, the immunophilins and the HSP 110 family. The low molecular weight family includes HSP 10, HSP 20, HSP 27 (Heme Oxygenase 1), HSP 32 and HSP 40. HSP 20 occurs in two complex sizes, dimers and multimers. It is related to stress proteins and occurs most abundantly in skeletal muscle and heart. HSP 20 is considerably shorter at the C-terminus and less polar than other small heat shock proteins.

REFERENCES

1. van de Klundert, FA., et al. 1998. The mammalian small heat-shock protein HSP 20 forms dimers and is a poor chaperone. *Eur. J. Biochem.* 258: 1014-1021.
2. Bukach, O.V., et al. 2004. Some properties of human small heat shock protein HSP 20 (HSPB 6). *Eur. J. Biochem.* 271: 291-302.
3. Gusev, N.B., et al. 2005. Structure, properties and probable physiological role of small heat shock protein with molecular mass 20 kD (HSP 20, HSP 6). *Biochemistry* 70: 629-637.
4. Batts, T.W., et al. 2005. Absence of force suppression in rabbit bladder correlates with low expression of heat shock protein 20. *BMC Physiol.* 5: 16.
5. Bukach, O.V., et al. 2005. Small heat shock protein with apparent molecular mass 20 kDa (HSP 20, HSP B6) is not a genuine Actin-binding protein. *J. Muscle Res. Cell Motil.* 26: 175-181.
6. Fontaine, J.M., et al. 2005. Interactions of HSP22 (HSPB8) with HSP 20, α B-crystallin and HSPB3. *Biochem. Biophys. Res. Commun.* 337: 1006-1011.

CHROMOSOMAL LOCATION

Genetic locus: Hspb6 (mouse) mapping to 7 B1.

SOURCE

HSP 20 (L-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HSP 20 of rat origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46140 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

HSP 20 (L-13) is recommended for detection of HSP 20 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HSP 20 siRNA (m): sc-45676, HSP 20 shRNA Plasmid (m): sc-45676-SH and HSP 20 shRNA (m) Lentiviral Particles: sc-45676-V.

Molecular Weight of HSP 20: 20 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **HSP 20 (HSP20-11): sc-51955**, our highly recommended monoclonal alternative to HSP 20 (L-13).