

p54/nrb (C-20): sc-46221

BACKGROUND

Found in both primary and transformed human cells, paraspeckles are discrete bodies in the interchromatin nucleoplasmic space which contain p54/nrb (nuclear RNA-binding protein) and at least two other RNA-binding proteins, paraspeckle protein 1 (PSP1) and paraspeckle protein 2 (PSP2). Paraspeckles often co-localize with splicing speckles, which are the site of splicing factor accumulation. Paraspeckle proteins, including p54/nrb, move dynamically between the nucleolus and paraspeckles and translocate to distinct caps in the nucleolar periphery when transcription is inhibited. Originally purified from HeLa cells, the nuclear p54/nrb has two RNA recognition motifs and shares extensive homology with both the human splicing factor PSF and *Drosophila* NONA/BJ6, which is required for normal vision and courtship. The shared domain between these proteins is termed a DBHS (*Drosophila* behavior, human splicing) domain and may play a role in regulating various pathways at the level of pre-mRNA splicing.

REFERENCES

- Dong, B., et al. 1993. Purification and cDNA cloning of HeLa cell p54/nrb, a nuclear protein with two RNA recognition motifs and extensive homology to human splicing factor PSF and *Drosophila* NONA/BJ6. *Nucleic Acids Res.* 21: 4085-4092.
- Brown, C.J., et al. 1997. Expression of genes from the human active and inactive X chromosomes. *Am. J. Hum. Genet.* 60: 1333-1343.
- Zhang, Z. and Carmichael, G.G. 2001. The fate of dsRNA in the nucleus: a p54/nrb-containing complex mediates the nuclear retention of promiscuously A-to-I edited RNAs. *Cell* 106: 465-475.
- Fox, A.H., et al. 2002. Paraspeckles: a novel nuclear domain. *Curr. Biol.* 12: 13-25.
- Shav-Tal, Y. and Zipori, D. 2002. PSF and p54(nrb)/NonO—multi-functional nuclear proteins. *FEBS Lett.* 531: 109-114.

CHROMOSOMAL LOCATION

Genetic locus: NONO (human) mapping to Xq13.1; Nono (mouse) mapping to X D.

SOURCE

p54/nrb (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of p54/nrb of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46221 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-46221 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p54/nrb (C-20) is recommended for detection of p54/nrb of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p54/nrb (C-20) is also recommended for detection of p54/nrb in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for p54/nrb siRNA (h): sc-38163, p54/nrb siRNA (m): sc-38164, p54/nrb shRNA Plasmid (h): sc-38163-SH, p54/nrb shRNA Plasmid (m): sc-38164-SH, p54/nrb shRNA (h) Lentiviral Particles: sc-38163-V and p54/nrb shRNA (m) Lentiviral Particles: sc-38164-V.

p54/nrb (C-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of p54/nrb: 54 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, MCF7 nuclear extract: sc-2149 or p54/nrb (h): 293T Lysate: sc-114638.

DATA



p54/nrb (C-20): sc-46221. Western blot analysis of p54/nrb expression in MCF7 (A), A-673 (B), PC-3 (C) and MOLT-4 (D) nuclear extracts.



p54/nrb (C-20): sc-46221. Western blot analysis of p54/nrb expression in non-transfected 293T: sc-117752 (A), human p54/nrb transfected 293T: sc-114638 (B) and HeLa (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **p54/nrb (F-5): sc-376804** or **p54/nrb (G-1): sc-376865**, our highly recommended monoclonal alternatives to p54/nrb (C-20).