

DCAMKL1 (N-14): sc-46312

BACKGROUND

Lissencephaly (smooth brain) is an abnormality of brain development characterized by incomplete neuronal migration and a smooth cerebral surface, manifesting as severe mental retardation. Genetic analysis has identified two proteins that are mutated in some cases of lissencephaly, designated lissencephaly-1 protein (LIS1, also designated platelet activating factor acetylhydrolase 45 kDa) and doublecortin. LIS1 displays sequence homology to β -subunits of heterotrimeric G proteins, and doublecortin contains a consensus Abl phosphorylation site. In addition, the DCAMKL1 (doublecortin-like and CAM kinase-like 1) protein shows homology to doublecortin. All three proteins are highly expressed in developing brain and may function together to regulate microtubules involved in neuronal migration. The DCAMKL1 protein encodes a functional kinase that is capable of phosphorylating myelin basic protein and itself, but its kinase activity does not appear to affect its microtubule polymerization activity.

CHROMOSOMAL LOCATION

Genetic locus: DCAMKL1 (human) mapping to 13q13.3; Dcamkl1 (mouse) mapping to 3 C.

SOURCE

DCAMKL1 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of DCAMKL1 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46312 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DCAMKL1 (N-14) is recommended for detection of DCAMKL1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DCAMKL1 (N-14) is also recommended for detection of DCAMKL1 in additional species, including equine and porcine.

Suitable for use as control antibody for DCAMKL1 siRNA (h): sc-45618, DCAMKL1 siRNA (m): sc-45619, DCAMKL1 shRNA Plasmid (h): sc-45618-SH, DCAMKL1 shRNA Plasmid (m): sc-45619-SH, DCAMKL1 shRNA (h) Lentiviral Particles: sc-45618-V and DCAMKL1 shRNA (m) Lentiviral Particles: sc-45619-V.


Molecular Weight of DCAMKL1: 82 kDa.

Positive Controls: mouse brain extract: sc-2253, rat brain extract: sc-2392 or NIH/3T3 whole cell lysate: sc-2210.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



DCAMKL1 (N-14): sc-46312. Western blot analysis of DCAMKL1 expression in rat brain (A) and mouse brain (B) tissue extracts.



DCAMKL1 (N-14): sc-46312. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **DCAMKL1 (D-3): sc-514684**, our highly recommended monoclonal alternative to DCAMKL1 (N-14).