

HIPK3 (C-14): sc-46358

BACKGROUND

The homeodomain-interacting protein kinase (HIPK) family includes three members, HIPK1, HIPK2, and HIPK3. Each family member contains a conserved protein kinase domain as well as a separate domain, which interacts with homeoproteins. HIPK2 appears to act as a corepressor of homeodomain transcription factors, such as NK-3. Also, HIPK2 is regulated by ubiquitin-like modification via the covalent binding of SUMO-1. Subsequently, it is directed to nuclear bodies *in vitro*.

REFERENCES

- Kim, Y.H., Choi, C.Y., Lee, S.J., Conti, M.A. and Kim, Y. 1998. Homeodomain-interacting protein kinases, a novel family of co-repressors for homeodomain transcription factors. *J. Biol. Chem.* 273: 25875-25879.
- Nupponen, N.N. and Visakorpi, T. 1999. Assignment of the protein kinase homolog of YAK1 (HIPK3) to human chromosome band 11p13 by *in situ* hybridization. *Cytogenet. Cell. Genet.* 87:102-103.
- Rochat-Steiner, V., Becker, K., Micheau, O., Schneider, P., Burns, K. and Tschopp, J. 2000. FIST/HIPK3: a Fas/FADD-interacting serine/threonine kinase that induces FADD phosphorylation and inhibits fas-mediated Jun NH₂-terminal kinase activation. *J. Exp. Med.* 192: 1165-1174.
- Curtin, J.F. and Cotter, T.G. 2004. JNK regulates HIPK3 expression and promotes resistance to Fas-mediated apoptosis in DU 145 prostate carcinoma cells. *J. Biol. Chem.* 279: 17090-17100.
- Venables, J.P., Bourgeois, C.F., Dalgliesh, C., Kister, L., Stevenin, J. and Elliott, D.J. 2005. Up-regulation of the ubiquitous alternative splicing factor Tra2 β causes inclusion of a germ cell-specific exon. *Hum. Mol. Genet.* 14: 2289-2303.
- Gresko, E., Moller, A., Roscic, A. and Schmitz, M.L. 2005. Covalent modification of human homeodomain interacting protein kinase 2 by SUMO-1 at lysine 25 affects its stability. *Biochem. Biophys. Res. Commun.* 329: 1293-1299.

CHROMOSOMAL LOCATION

Genetic locus: HIPK3 (human) mapping to 11p13; Hipk3 (mouse) mapping to 2 E2.

SOURCE

HIPK3 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of HIPK3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46358 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

HIPK3 (C-14) is recommended for detection of HIPK3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HIPK3 (C-14) is also recommended for detection of HIPK3 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for HIPK3 siRNA (h): sc-45654, HIPK3 siRNA (m): sc-45655, HIPK3 shRNA Plasmid (h): sc-45654-SH, HIPK3 shRNA Plasmid (m): sc-45655-SH, HIPK3 shRNA (h) Lentiviral Particles: sc-45654-V and HIPK3 shRNA (m) Lentiviral Particles: sc-45655-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.