

HoxD9 (N-15): sc-46367

BACKGROUND

The Hox proteins play a role in patterns of embryonic development and cellular differentiation by regulating downstream target genes. *In vivo*, the HoxD9 protein interacts with the autoregulatory and cross-regulatory enhancers of the murine HoxB1 and human HoxD9 genes. Specifically, the HoxD9 protein interacts with the human control region (HCR) of the HoxD9 gene, thus inducing transcription of the HoxD9 promoter. HoxD9 may be a multifunctional transcriptional regulator, as it contains different activation domains. Activation of HoxD9 depends on the structure of the target regulatory element, and results in differential cofactor interaction. The HoxD9 protein is expressed in the early stages of mouse joint development, primarily in the articular cartilage. HoxD9 transcripts are also detected in the synovial tissue of arthritic mice, but not in that of normal mice, suggesting that HoxD9 may have a role in the pathology of arthritis. Furthermore, the HoxD9 protein is highly expressed in the synoviocytes of patients with rheumatoid arthritis (RA), but not in osteoarthritis patients. The human HoxD9 protein is also differentially expressed in the human cervical cancer cell line HeLa, but is not expressed in the normal cervix and may thus play a role in tumorigenesis.

CHROMOSOMAL LOCATION

Genetic locus: HOXD9 (human) mapping to 2q31.1; Hoxd9 (mouse) mapping to 2 C3.

SOURCE

HoxD9 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of HoxD9 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-46367 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-46367 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HoxD9 (N-15) is recommended for detection of HoxD9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HoxD9 siRNA (h): sc-35585, HoxD9 siRNA (m): sc-35586, HoxD9 shRNA Plasmid (h): sc-35585-SH, HoxD9 shRNA Plasmid (m): sc-35586-SH, HoxD9 shRNA (h) Lentiviral Particles: sc-35585-V and HoxD9 shRNA (m) Lentiviral Particles: sc-35586-V.

HoxD9 (N-15) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of HoxD9: 28-41 kDa.

Positive Controls: HoxD9 (h): 293T Lysate: sc-115615, FHs 173We cell lysate: sc-2417 or HoxD9 (m): 293T Lysate: sc-126973.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



HoxD9 (N-15): sc-46367. Western blot analysis of HoxD9 expression in non-transfected: sc-117752 (A) and human HoxD9 transfected: sc-115615 (B) 293T whole cell lysates.



HoxD9 (N-15): sc-46367. Western blot analysis of HoxD9 expression in non-transfected: sc-117752 (A) and mouse HoxD9 transfected: sc-126973 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **HoxD9 (H-2): sc-137134** or **HoxD9 (B-9): sc-365717**, our highly recommended monoclonal alternatives to HoxD9 (N-15).