

SULT2B1 (N-15): sc-46542

BACKGROUND

The soluble sulfotransferases contribute to the elimination of xenobiotics, the activation of procarcinogens, and the regulation of hormones. Members of the three groups comprising this superfamily show selectivity to certain substrate compounds. SULT1 sulfotransferases exhibit N-sulfating activities of carcinogenic heterocyclic amines, and are selective toward phenols, whereas SULT2 enzymes prefer hydroxysteroids and SULT3 family members are selective for N-substituted aryl and alicyclic compounds. SULT2A1 catalyzes the sulfonation of procarcinogen xenobiotics, hydroxysteroids and bile acids, and is highly expressed in adrenal and liver tissues. SULT2A1 plays a role in hepatic cholesterol homeostasis. SULT2B1 consists of two isoforms, SULT2B1a and SULT2B1b, which are transcribed from the same gene by alternative splicing of their first exons. Both isoforms are highly selective for the sulphation of 3 β -hydroxysteroids, such as pregnenolone, epiandrosterone, DHEA and androstenediol. SULT2B1b is expressed in prostate, skin, placenta and lung tissues.

CHROMOSOMAL LOCATION

Genetic locus: SULT2B1 (human) mapping to 19q13.33; Sult2b1 (mouse) mapping to 7 B4.

SOURCE

SULT2B1 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SULT2B1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46542 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SULT2B1 (N-15) is recommended for detection of SULT2B1 isoforms 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with SULT2A1.

SULT2B1 (N-15) is also recommended for detection of SULT2B1 isoforms 1 and 2 in additional species, including canine.

Suitable for use as control antibody for SULT2B1 siRNA (h): sc-44399, SULT2B1 siRNA (m): sc-44400, SULT2B1 shRNA Plasmid (h): sc-44399-SH, SULT2B1 shRNA Plasmid (m): sc-44400-SH, SULT2B1 shRNA (h) Lentiviral Particles: sc-44399-V and SULT2B1 shRNA (m) Lentiviral Particles: sc-44400-V.

Molecular Weight of mouse and rat SULT2B1: 30 kDa.


Molecular Weight of human SULT2B1: 41 kDa.

Positive Controls: rat kidney extract: sc-2394, mouse small intestine extract: sc-364252 or SULT2B1 (h): 293T Lysate: sc-113758.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SULT2B1 (N-15): sc-46542. Western blot analysis of SULT2B1 expression in non-transfected: sc-117752 (A) and human SULT2B1 transfected: sc-113758 (B) 293T whole cell lysates.



SULT2B1 (N-15): sc-46542. Western blot analysis of SULT2B1 expression in mouse small intestine tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SULT2B1 (A-3): sc-166423** or **SULT2B1 (C-1): sc-166346**, our highly recommended monoclonal alternatives to SULT2B1 (N-15).