

Transketolase (N-19): sc-46554

BACKGROUND

Transketolase (TK or TKT), a member of the Transketolase family, is a multi-functional protein that plays a role in diabetes, cancer, Alzheimer's disease and Wernicke-Korsakoff's syndrome, a latent genetic neurological disorder. Transketolase is also important for the prevention of hyperglycemia-induced vascular damage. Transketolase is a crucial protein in the pentose phosphate pathway (PPP), where it catalyzes several reactions. In combination with Transaldolase, Transketolase functions as a link between glycolysis and the non-oxidative part of the PPP, allowing the cell to adapt to varying metabolic conditions in response to environmental changes. Transketolase activity is detected in small intestine epithelia, liver parenchyma, tongue, cornea and trachea. It is also expressed in the proximal tubules of kidney and in ganglion cells in medulla of the adrenal gland.

CHROMOSOMAL LOCATION

Genetic locus: TKT (human) mapping to 3p21.1; Tkt (mouse) mapping to 14 B.

SOURCE

Transketolase (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Transketolase of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46554 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Transketolase (N-19) is recommended for detection of Transketolase of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Transketolase (N-19) is also recommended for detection of Transketolase in additional species, including bovine, porcine and avian.

Suitable for use as control antibody for Transketolase siRNA (h): sc-45591, Transketolase siRNA (m): sc-45592, Transketolase shRNA Plasmid (h): sc-45591-SH, Transketolase shRNA Plasmid (m): sc-45592-SH, Transketolase shRNA (h) Lentiviral Particles: sc-45591-V and Transketolase shRNA (m) Lentiviral Particles: sc-45592-V.


Molecular Weight of Transketolase: 78 kDa.

Positive Controls: KNRK whole cell lysate: sc-2214, mouse eye extract: sc-364241 or rat liver extract: sc-2395.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Transketolase (N-19): sc-46554. Western blot analysis of Transketolase expression in KNRK whole cell lysate (A) and rat kidney (B), rat liver (C) and mouse eye (D) tissue extracts.



Transketolase (N-19): sc-46554. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human salivary gland tissue showing nuclear and cytoplasmic staining of glandular cells (B).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Transketolase (H-7): sc-390179**, our highly recommended monoclonal alternative to Transketolase (N-19).