



Epigen (N-19): sc-46942

BACKGROUND

Epigen is an epithelial mitogen that belongs to the EGF ligand family for ErbB receptors. Comprising 152 amino acids, the Epigen protein contains two hydrophobic regions corresponding to a putative signal sequence and transmembrane domain, which flank a core of amino acids encompassing six cysteine residues and two putative N-linked glycosylation sites. In epithelial cells, Epigen stimulates the phosphorylation of ErbB-1 and mitogen-activated protein kinases and also activates a reporter gene containing enhancer sequences present in the c-Fos promoter. Epigen also stimulates the proliferation of HaCa T cells. With its ability to promote the growth of epithelial cells, Epigen may constitute a novel molecular target for wound-healing therapy.

REFERENCES

1. Strachan, L., Murison, J.G., Prestidge, R.L., Sleeman, M.A., Watson, J.D. and Kumble, K.D. 2001. Cloning and biological activity of Epigen, a novel member of the EP growth-factor superfamily. *J. Biol. Chem.* 276: 18265-18271.
2. Lee, D., Pearsall, R.S., Das, S., Dey, S.K., Godfrey, V.L. and Threadgill, D.W. 2004. Epiregulin is not essential for development of intestinal tumors but is required for protection from intestinal damage. *Mol. Cell. Biol.* 24: 8907-8916.
3. Kochupurakkal, B.S., Harari, D., Di-Segni, A., Maik-Rachline, G., Lyass, L., Gur, G., Kerber, G., Citri, A., Lavi, S., Eilam, R., Chalifa-Caspi, V., Eshhar, Z., Pikarsky, E., Pinkas-Kramarski, R., Bacus, S.S. and Yarden, Y. 2005. Epigen, the last ligand of ErbB receptors, reveals intricate relationships between affinity and mitogenicity. *J. Biol. Chem.* 280: 8503-8512.
4. Katoh, Y. and Katoh, M. 2006. Canonical Wnt-signaling pathway and human AREG. *Int. J. Mol. Med.* 17: 1163-1166.
5. Amsellem-Ouazana, D., Bièche, I., Tozlu, S., Botto, H., Debre, B. and Lidereau, R. 2006. Gene expression profiling of ErbB receptors and ligands in human cell carcinoma of the bladder. *J. Urol.* 175: 1127-1132.

CHROMOSOMAL LOCATION

Genetic locus: EPGN (human) mapping to 4q13.3; Epgn (mouse) mapping to 5 E1.

SOURCE

Epigen (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Epigen of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-46942 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Epigen (N-19) is recommended for detection of Epigen of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Epigen siRNA (m): sc-60592; and as shRNA Plasmid control antibody for Epigen shRNA Plasmid (m): sc-60592-SH.

Molecular Weight of Epigen: 15 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.