

GPT (N-20): sc-47020

BACKGROUND

The glutamate pyruvate transaminases GPT (or GPT1) and GPT2, also designated alanine aminotransferases (ALT1 and ALT2), respectively, catalyze the reversible transamination between alanine and 2-oxoglutarate to form pyruvate and Glutamate. Subsequently, they play a key role in the intermediary metabolism of glucose and amino acids. GPT and GPT2 share significant sequence homology, but differ in their expression patterns. GPT exhibits high expression in kidney, liver and heart, whereas GPT2 expression is high in muscle, fat and kidney. GPT is widely used as an index of liver integrity or hepatocellular damage in clinical settings.

REFERENCES

1. Sohocki, M.M., et al. 1997. Human glutamate pyruvate transaminase (GPT): localization to 8q24.3, cDNA and genomic sequences and polymorphic sites. *Genomics* 40: 247-252.
2. Yang, R.Z., et al. 2002. cDNA cloning, genomic structure, chromosomal mapping and functional expression of a novel human alanine aminotransferase. *Genomics* 79: 445-450.

CHROMOSOMAL LOCATION

Genetic locus: GPT (human) mapping to 8q24.3; Gpt (mouse) mapping to 15 D3.

SOURCE

GPT (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of GPT of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47020 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GPT (N-20) is recommended for detection of GPT of human and, to a lesser extent, mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPT siRNA (h): sc-60753, GPT siRNA (m): sc-60754, GPT shRNA Plasmid (h): sc-60753-SH, GPT shRNA Plasmid (m): sc-60754-SH, GPT shRNA (h) Lentiviral Particles: sc-60753-V and GPT shRNA (m) Lentiviral Particles: sc-60754-V.


Molecular Weight of GPT: 48 kDa.

Positive Controls: GPT (h): 293 Lysate: sc-111013 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



GPT (N-20): sc-47020. Western blot analysis of GPT expression in non-transfected: sc-110760 (A) and human GPT transfected: sc-111013 (B) 293 whole cell lysates.



GPT (N-20): sc-47020. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing cytoplasmic and nuclear staining of hepatocytes (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing nuclear and cytoplasmic staining of hepatocytes. Kindly provided by The Swedish Human Protein Atlas (HPA) program (B).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **GPT (E-3): sc-374501** or **GPT (G-9): sc-271089**, our highly recommended monoclonal alternatives to GPT (N-20).