

IL-23R (C-13): sc-47060

BACKGROUND

The p19 protein shares sequence similarity with IL-6 subfamily members and is distantly related to the p35 subunit of IL-12. p19 shows no biological activity by itself; instead, it combines with the p40 subunit of IL-12 to form a biologically active, composite cytokine, IL-23. IL-23 shares some *in vivo* functions with IL-12, including the activation of the transcription factor Stat4. Also similar to IL-12, human IL-23 stimulates IFN- γ production and proliferation in PHA blast T cells, as well as in CD45RO (memory) T cells. Ubiquitous transgenic expression of the IL-23 subunit p19 induces multiorgan inflammation, runting, infertility and premature death. The receptors for each appear to share one subunit, but also have at least one distinct subunit. Activated dendritic cells secrete detectable levels of this heterodimeric complex and IL-23 binds to IL-12R β 1 and IL-23R. At least six spliced isoforms of IL-23R (IL-23R1 to 6) can be generated through alternative splicing. The IL-23R2 and/or IL-23R4 variants are predominantly detected in certain human lung carcinomas and may contribute to their pathogenesis.

REFERENCES

1. Oppmann, B., et al. 2000. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 13: 715-725.
2. Wiekowski, M.T., et al. 2001. Ubiquitous transgenic expression of the IL-23 subunit p19 induces multiorgan inflammation, runting, infertility and premature death. *J. Immunol.* 166: 7563-7570.
3. Frucht, D.M. 2002. IL-23: a cytokine that acts on memory T cells. *Sci. STKE* 2002: PE1.
4. Cooper, A.M., et al. 2002. Mice lacking bioactive IL-12 can generate protective, antigen-specific cellular responses to mycobacterial infection only if the IL-12 p40 subunit is present. *J. Immunol.* 168: 1322-1327.
5. Watford, WT., et al. 2004. Signaling by IL-12 and IL-23 and the immunoregulatory roles of STAT4. *Immunol. Rev.* 202: 139-156.
6. Vanden Eijnden, S., et al. 2005. IL-23 upregulates IL-10 and induces IL-17 synthesis by polyclonally activated naive T cells in human. *Eur. J. Immunol.* 35: 469-475.
7. Zhang, X.Y., et al. 2006. Identification and expression analysis of alternatively spliced isoforms of human interleukin-23 receptor gene in normal lymphoid cells and selected tumor cells. *Immunogenetics* 57: 934-943.

CHROMOSOMAL LOCATION

Genetic locus: IL23R (human) mapping to 1p31.2; IL23a (mouse) mapping to 10 D3.

SOURCE

IL-23R (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of IL-23R of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47060 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

IL-23R (C-13) is recommended for detection of IL-23R of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-23 siRNA (h): sc-43860, IL-23 siRNA (m): sc-60028, IL-23 shRNA Plasmid (h): sc-43860-SH, IL-23 shRNA Plasmid (m): sc-60028-SH, IL-23 shRNA (h) Lentiviral Particles: sc-43860-V and IL-23 shRNA (m) Lentiviral Particles: sc-60028-V.

Molecular Weight of IL-23R: 69 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **IL-23R (3D7): sc-293485**, our highly recommended monoclonal alternative to IL-23R (C-13).